PATENT AND TRADEMARK OFFICE	12 Rec'd PCT/PTO 1 0 APR 20
AND ADADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER
UNITED STATES	A33153-PCT USA
E (DO/EO/US)	U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR
R 35 U.S.C. 371	09/529239
ONAL FILING DATE	PRIORITY DATE CLAIMED
9 October 1998	10 October 1997
IES s S.; FREYSSINET, Georg	ges; and PEREZ, Pascal
d/Elected Office (DO/EO/US) t	he following items and other information:
ing a filing under 35 U.S.C. 371	
ssion of items concerning a fili	
-	C. 371(f)) at any time rather than delay 371(b) and PCT Articles 22 and 39(1).
y Examination was made by the	e 19th month from the earliest claimed priority date.
ed (35 U.S.C. 371 (c) (2))	
y if not transmitted by the Inter	rnational Bureau).
onal Bureau.	
s filed in the United States Rec	eiving Office (RO/US).
into English (35 U.S.C. 371(c)(2)).
CT/ISA/210).	
l Application under PCT Article	e 19 (35 U.S.C. 371 (c)(3))
aly if not transmitted by the Inte	ernational Bureau).
tional Bureau.	
me limit for making such amend	dments has NOT expired.
made.	
under PCT Article 19 (35 U.S.	C. 371(c)(3)).
J.S.C. 371 (c)(4)).	
ination Report (PCT/IPEA/409	
al Preliminary Examination Re	port under PCT Article 36
nation included:	
37 CFR 1.97 and 1.98.	
parate cover sheet in complianc	e with 37 CFR 3.28 and 3.31 is included.
amendment.	
letter.	
m PCT/IB/308; Form PCT/II	PEA/ 416; a postcard, and a check in the amount
10	orm PCT/IB/308; Form PCT/II

The line good from all their role had not not had live it

- 11. A DNA molecule according to claim 10 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 12. A DNA molecule according to claim 10 further comprising a regulation element capable of causing overexpression of said polypeptide in a cell of said plant.
- 5 13. A chimeric gene comprising:
- a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant; and

at least one regulation element capable of functioning in a plant cell.

- 14. A chimeric gene according to claim 13 wherein said regulation element is selected from constitutive, inducible, tissue type specific and cell type specific promoters.
- 15. A chimeric gene according to claim 13 comprising a DNA sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant, wherein said regulation element is capable of causing overexpression of said polypeptide in a cell of said plant.
- 16. A chimeric gene according to claim 13 wherein said regulation element is selected from the group consisting of 35S, NOS, PR1a, AoPR1 and DMC1.
 - 17. A plasmid or vector comprising a chimeric gene according to any one of claims 13-16.
 - 18. A plant cell stably transformed, transfected or electroporated with a plasmid or vector according to claim 17.
- 19. A plant comprising a cell according to claim 18.
 - 20. A plant according to claim 19 selected from plants of the families Brassicaceae, Poaceae, Solanaceae, Asteraceae, Malvaceae, Fabaceae, Linaceae, Canabinaceae, Dauaceae and Cucurbitaceae.
- 21. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a DNA molecule according to any one of claims 1-3 or 7-12 and causing said DNA sequence to express said polynucleotide or said polypeptide.
 - 22. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a chimeric gene

422 Rec'd PCT/PTO 1 0 APR 2000

	NO. (IF KNOWN, SEE 37 CFR	INTERNATIONAL APPLICAT	ON NO.		ATTORNEY'S	DOCKET NUMBER
<u> </u>	<u> 529239</u>	PCT/EP98/069	77		A33153-	PCT USA
21. The foll	owing fees are submitted:.				CALCULATIONS	PTO USE ONLY
	L FEE (37 CFR 1.492 (a) (1) -					
international	national preliminary examination search fee (37 CFR 1.445(a)(2) onal Search Report not prepared	paid to USPTO	\$9	70.00		
☑ International USPTO but I	preliminary examination fee (37 nternation Search Report prepar	CFR 1.482) not paid to ed by the EPO or JPO	\$8	40.00		
but internation	preliminary examination fee (37 onal search fee (37 CFR 1.445(a)	(2)) paid to USPTO		90.00		
but all claims	preliminary examination fee pair did not satisfy provisions of PC	T Article 33(1)-(4)	\$6	70.00		
☐ International and all claims	preliminary examination fee pais satisfied provisions of PCT Ar		•	96.00		
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		ATE BASIC FEE AM		=	\$840.00	
months from the earl	0 for furnishing the oath or declining the claimed priority date (37 C	FR 1.492 (e)).			\$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RAT			
Total claims	64 - 20 =	44	x \$18		\$792.00	
Independent claims	9 - 3=	6	x \$78	.00	\$468.00	
Multiple Dependent	Claims (check if applicable).	ABOVE CALCULAT			\$260.00 \$2,360.00	
Peduction of 1/2 for	filing by small entity, if applica			T	\$2,500.00	
must also be filed ()	Note 37 CFR 1.9, 1.27, 1.28) (ch	eck if applicable).			\$0.00	
		SUB	<u>TOTAI</u>	, =	\$2,360.00	
Processing fee of \$1.	30.00 for furnishing the English liest claimed priority date (37 C	translation later than \Box 2	0 🗆	30 +	Ф0.00	
inolitis from the ear	mest claimed priority date (37 C		LEDD		\$0.00	
		TOTAL NATIONA		=	\$2,360.00	
Fee for recording the accompanied by an a	e enclosed assignment (37 CFR appropriate cover sheet (37 CFR	1.21(h)). The assignment must 3.28, 3.31) (check if applicab	be le).		\$0.00	
		TOTAL FEES ENCI	LOSED	=	\$2,360.00	
re re					Amount to be: refunded	\$
					charged	\$
🛚 🛚 A check in t	the amount of \$2,360.00	to cover the above fees is en	closed.			
1	ge my Deposit Account No.	in the amount o	f		to cover the above	ve fees.
A duplicate	e copy of this sheet is enclosed.					
1	ssioner is hereby authorized to of Account No. 02-4377	charge any fees which may be re A duplicate copy of this sheet is	•		ny overpayment	
NOTE: Where an	appropriate time limit under 3	7 CFR 1.494 or 1.495 has not	been met,		ion to revive (37 CF	R
	st be filed and granted to resto	re the application to pending	status.)		
SEND ALL CORRE	SPONDENCE TO:		KO	1/11/	W. Sul	ll
Rochelle K. Seide			SIGNA	TURE		
Baker Botts LLP			Roche	lle K. S	leide	
30 Rockefeller Pla New York, NY 10			NAME			
US	·		32,300			
					ON NUMBER	
			10 Apı	:i1 2000	1	
			DATE	2000		
			DATE			

ţ

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant

Doutriaux, et al.

Serial No.

Not Yet Assigned

Examiner:

Filed

April 10, 2000

Group Art Unit:

For

METHODS FOR OBTAINING PLANT VARIETIES

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents

Washington, D.C. 20231

Sir:

Prior to the examination of the present application, please make the

following amendments.

IN THE CLAIMS:

Please make the following amendments:

Please renumber the second Claim "25" as --26--.

Please renumber Claim "26" as --27--.

Please renumber Claim "27" as --28--.

Please renumber Claim "28" as --29--; and in the first line thereof, change

and the second of the second

"27" to --28--.

NY02:257264.1

PATENT

Please renumber Claim "29" as --30--; and in the first line thereof, change "28" to --29--.

Please renumber Claim "31" as --32--; and in the first line thereof, change "27" to --28--.

Please renumber Claim "32" as --33--; and in the first line thereof, change "31" to --32--.

Please renumber Claim "33" as --34--.

Please renumber Claim "34" as --35--.

Please renumber Claim "35" as --36--.

IN THE ABSTRACT

After the Claims, please insert the following Abstract:

--An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.--

REMARKS

The present amendment is necessitated to eliminate the duplicate numbering of Claim 25, and to clarify the claim numbering and dependencies of the

PATENT

remaining claims. Also, an Abstract is provided. No new matter has been added.

Respectfully submitted,

Rochelle K. Seide

Patent Office Reg. No. 32,300

Attorneys for Applicants (212) 408-2626

Methods for Obtaining Plant Varieties

1

TECHNICAL FIELD

The present invention relates to nucleotide sequences which encode polypeptides involved in the DNA mismatch repair systems of plants, and to the polypeptides encoded 5 by those nucleotide sequences. The invention also relates to nucleotide sequences and polypeptide sequences for use in altering the DNA mismatch repair system in plants. The invention also relates to a process for altering the DNA mismatch repair system of a plant cell, to a process for increasing genetic variations in plants and to processes for obtaining plants having a desired characteristic.

BACKGROUND OF THE INVENTION

Plant breeding essentially relies on and makes use of genetic variation which occurs naturally within and between members of a family, a genus, a species or a subspecies. Another source of genetic variation is the introduction of genes from other organisms which may or may not be related to the host plant.

Allelic loci or non-allelic genes which constitute or contribute to desired quantitative (e.g. growth performance, yield, etc.) or qualitative (e.g. deposition, content and composition of seed storage products; pathogen resistance genes; etc.) traits that are absent, incomplete or inefficient in a species or subspecies of interest are typically introduced by the plant breeder from other species or subspecies, or de novo. This 20 introduction is often done by crossing, provided that the species to be crossed are sexually compatible. Other means of introducing genomes, individual chromosomes or genes into plant cells or plants are well known in the art. They include cell fusion, chemically aided transfection (Schocher et al., 1986, Biotechnology 4: 1093) and ballistic (McCabe et al., 1988, Biotechnology 6: 923), microinjection (Neuhaus et al., 1987, TAG 75: 30), 25 electroporation of protoplasts (Chupeau et al., 1989, Biotechnology 7: 53) or microbial transformation methods such as Agrobacterium mediated transformation (Horsch et al., 1985, Science 227: 1229; Hiei et al., 1996, Biotechnology 14: 745).

However, when a foreign genome, chromosome or gene is introduced into a plant, it will often segregate in subsequent generations from the genome of the recipient plant or 30 plant cell during mitotic and meiotic cell divisions and, in consequence, become lost from the host plant or plant cell into which it had been introduced. Occasionally, however, the introduced genome, chromosome or gene physically combines entirely or in part with the genome, chromosome or gene of the host plant or plant cell in a process which is called recombination.

Recombination involves the exchange of covalent linkages between DNA molecules 35 in regions of identical or similar sequence. It is referred to here as homologous recombination if donor and recipient DNA are identical or nearly identical (at least 99%

in the second second second

And the state of t

4.

10

15

L.j

4

base sequence identity), and as homeologous recombination if donor and recipient DNA are not identical but are similar (less than 99% base sequence identity).

The ability of two genomes, chromosomes or genes to recombine is known to depend largely on the evolutionary relation between them and thus on the degree of sequence similarity between the two DNA molecules. Whereas homologous recombination is frequently observed during mitosis and meiosis, homeologous recombination is rarely or never seen.

From a breeder's perspective, the limits within which homologous recombination occurs, therefore, define a genetic barrier between species, varieties or lines, in contrast to homeologous recombination which can break this barrier. Homeologous recombination is thus of great importance for plant breeding. Accordingly there is a need for a process for enhancing the frequency of homeologous recombination in plants. In particular, there is a need for a process of increasing homeologous recombination to significantly shorten the length of breeding programs by reducing the number of crosses required to obtain an otherwise rare recombination event.

At least in Escherichia coli, homologous and homeologous recombination are known to share a common pathway that requires among others the proteins RecA, RecB, RecC. RecD and makes use of the SOS induced RuvA and RuvB, respectively. It has been suggested that mating induced recombination follows the Double-Strand Break Repair 20 model (Szostak et al., 1983, Cell 33, 25-35), which is widely used to describe genetic recombination in eukaryotes. Following the alignment of homologous or homeologous DNA double helices the RecA protein mediates an exchange of a single DNA strand from the donor helix to the aligned recipient DNA helix. The incoming strand screens the recipient helix for sequence complementarity, seeking to form a heteroduplex by hydrogen 25 bonding the complementary strand. The displaced homologous or homeologous strand of the recipient helix is guided into the donor helix where it base pairs with its counterpart strand to form a second heteroduplex. The resulting branch point then migrates along the aligned chromosomes thereby elongating and thus stabilising the initial heteroduplexes. Single stranded gaps (if present) are closed by DNA synthesis. The strand cross overs 30 (Holliday junction) are eventually resolved enzymatically to yield the recombination products.

Although in wild type *E. coli* homologous and homeologous recombination are thus mechanistically similar if not identical, homologous recombination in conjugational crosses *E. coli* x *E. coli* occurs five orders of magnitude more frequently than homeologous recombination in conjugational crosses *E. coli* x *S. typhimurium* (Matic et al. 1995; Cell 80, 507-515). The imbalance in favour of homologous recombination was shown to be caused largely by the bacterial MisMatch Repair (MMR) system since its

The second second second second

WO 99/19492 PCT/EP98/06977

inactivation increased the frequency of homeologous recombination in *E. coli* up to 1000 fold (Rayssiguier et al. 1989, Nature 342, 396-401).

In E. coli, the MMR system (reviewed by Modrich 1991, Annual Rev Genetics 25, 229-253) is composed of only three proteins known as MutS, MutL and MutH. MutS recognizes and binds to base pair mismatches. MutL then forms a stable complex with mismatch bound MutS. This protein complex now activates the MutH intrinsic single stranded endonuclease which nicks the strand containing the misplaced base and thereby prepares the template for DNA repair enzymes.

During recombination, MMR components inhibit homeologous recombination. In vitro experiments demonstrated that MutS in complex with MutL binds to mismatches at the recombination branch point and physically blocks RecA mediated strand exchange and heteroduplex formation (Worth et al., 1994; PNAS 91, 3238-3241). Interestingly, the SOS dependent RuvAB mediated branch migration is insensitive to MutS/MutL, explaining the observed slight increase in SOS dependent homeologous recombination.

15 Homeologous mating even induces the SOS response, thereby taking advantage of RuvAB induction (Matic et al. 1995, Cell 80, 507-515).

The MMR system thus appears to be a genetic guardian over genome stability in *E. coli*. In this role it essentially determines the extent to which genetic isolation, that is, speciation, occurs. The diminished sensitivity of the SOS system to MMR, however, allows (within limits) for rapid genomic changes at times of stress, providing the means for fast adaptation to altered environmental conditions and thus contributing to intraspecies genetic variation and species evolution.

The important role of MMR in preserving genomic integrity has been established also in certain eukaryotes. In its efficiency, the human MMR, for example, may even counteract potential gene therapy tools such as triple-helix forming oligonucleotides including RNA-DNA hybrid molecules (Havre et al., 1993, J. Virology 67: 7234-7331; Wang et al., 1995, Mol. Cell. Biol. 15: 1759-1768; Kotani et al., 1996, Mol. Gen. Genetics 250: 626-634; Cole-Strauss et al., 1996, Science 273: 1387-1389). Such oligonucleotides are designed to introduce single base changes into selected DNA target sequences in order to inactivate for example cancer genes or to restore their normal function. The resulting base mismatches however are recognised by the mismatch repair system which then directs removal of the mismatched base, thereby reducing the efficiency of oligonucleotide induced site-specific mutagenesis.

To date, two families of related genes, homologous to the bacterial *MutS* and *MutL* genes have been identified or isolated in yeast and mammals (recent reviews by Arnheim and Shibata, 1997, Curr. Opinion Genet. Dev. 7, 364-370; Modrich and Lahue, 1996, Annual Rev. Biochem. 65, 101-133; Umar and Kunkel, 1996, Eur. J. Biochem. 238, 297-307). Biochemical and genetic analysis indicated that eukaryotic MutS homologs (MSH)

N ...

25

and MutL homologs (MLH, PMS), respectively, fulfil similar protein functions as their bacterial counterparts. Their relative abundance, however, could reflect different mismatch specificity and/or specialisation for different tissues or organelles or developmental processes such as mitotic versus meiotic recombination.

To date, six different genes homologous to *MutS* have been isolated in yeast (yMSH), and their homologs have been found in mouse (mMSH) and human (hMSH), respectively. Encoded proteins yMSH2, yMSH3 and yMSH6 appear to be the main MutS homologs involved in MMR during mitosis and meiosis in yeast, where the complementary proteins MSH3 and MSH6 alternatively associate with MSH2 to recognise different mismatch substrates (Masischky et al., 1996, Genes Dev. 10, 407-420). Similar protein interactions have been demonstrated for the human homologs hMSH2, hMSH3 and hMSH6 (Acharya et al., 1996, PNAS 93, 13629-13634).

MutL homologs (MLH and PMS), recently reviewed by Modrich and Lahue (1996, Annual Rev. Biochem. 65, 101-133) have so far been found in yeast (yMLH1 and yPMS1), mouse (mPMS2) and human (hMLH1, hPMS1 and hPMS2). The hPMS2 is a member of a family of at least 7 genes (Horii et al., 1994, Biochem. Biophys. Res. Commun. 204, 1257-1264) and its gene product is most closely related to yPMS1. Prolla et al. (1994, Science 265, 1091-1093) presented evidence for yPMS1 and yMLH1 to physically associate with each other and, together, to interact with the MutS homolog yMSH2 to form a ternary complex involved in mismatch substrate binding.

However, while medical interest in mismatch repair has prompted extensive research on MMR in bacteria, yeast and mammals, MMR genes have not been isolated from higher plants prior to the present invention and no attempts to adjust the plant MMR to plant breeding needs have been reported.

SUMMARY OF THE INVENTION

According to a first embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant. In one form of this embodiment, the invention provides an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human. More particularly, the invention provides polynucleotide sequences encoding polypeptides which are homologous to the mismatch repair polypeptides MSH3 and MSH6 of Saccharomyces cerevisiae. Still more particularly, the invention provides the coding sequences of the genes AtMSH3 and AtMSH6 of Arabidopsis thaliana. as defined hereinbelow, and polynucleotide sequences encoding polypeptides which are homologous to polypeptides encoded by AtMSH3 and AtMSH6.

adilla a cara

•

According to a second embodiment of the invention, there is provided an isolated and purified polypeptide functionally involved in the DNA mismatch repair system of a plant, for example a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human such as a polypeptide encoded by the genes AtMSH3 or AtMSH6 of Arabidopsis thaliana, as defined hereinbelow.

According to a third embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

According to a fourth embodiment of the invention there is provided a chimeric gene comprising a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant: together with at least one regulation element capable of functioning in a plant cell. Examples of such regulation elements include constitutive, inducible, tissue type specific and cell type specific promoters such as 35S, NOS, PR1a, AoPR1 and DMC1. Typically, a chimeric gene of the fourth embodiment will also include at least one terminator sequence, more typically exactly one terminator sequence.

In the third and fourth embodiments, said interference, by said polynucleotide sequence, with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair peptide of a yeast or a human typically occurs by hybridisation or by co-suppression.

According to a fifth embodiment of the invention there is provided a plasmid or vector comprising a chimeric gene of the fourth embodiment. A vector of the fifth 30 embodiment may be, for example, a viral vector or a bacterial vector.

According to a sixth embodiment of the invention, there is provided a plant cell stably transformed, transfected or electroporated with a plasmid or vector of the fifth embodiment.

According to seventh embodiment of the invention, there is provided a plant comprising a cell of the sixth embodiment.

According to an eighth embodiment of the invention, there is provided a process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising

The state of the state of the state of

1.1

transforming or transfecting said plant cell with a DNA sequence of the third embodiment or a chimeric gene of the fourth embodiment or a plasmid or vector of the fifth embodiment, and causing said DNA sequence to express said polynucleotide or said polypeptide.

According to a ninth embodiment of the invention, there is provided a process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred. For example, homeologous recombination may be evidenced by new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait.

According to a tenth embodiment of the invention there is provided a process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.

In a preferred form of the ninth and tenth embodiments of the invention, the step of altering the mismatch repair system comprises introducing into said hybrid plant, plant. cell or cells a chimeric gene of the fourth embodiment and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant or cells.

In other embodiments, the invention provides (a) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule of the first embodiment; (b) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO: 18 and (c) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30; with the proviso that the oligonucleotide of (a), (b) and (c) is other than SEQ ID NO:1 or SEQ ID NO:2.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a diagrammatic representation of the primer sequences used to 35 isolate AtMSH3.

Figure 2 is a plasmid map of clone 52, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH3.

Figure 3 is a plasmid map of clone 13, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for *AtMSH3*.

Figure 4 is a sequence listing of the coding sequence of AtMSH3, together with a deduced sequence of the encoded polypeptide.

Figure 5 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH3 protein.

Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6.

Figure 7 is a plasmid map of clone 43, showing restriction enzyme cleavage sites in to the 5' half of the full-length cDNA for *AtMSH6*.

Figure 8 is a plasmid map of clone 62, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for *AtMSH6*.

Figure 9 is a sequence listing of the coding sequence of *AtMSH6*, together with a deduced sequence of the encoded polypeptide.

Figure 10 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH6 protein.

Figure 11 is a genomic sequence listing of AtMSH6.

Figure 12 is a plasmid map of plasmid pPF13.

Figure 13 is a plasmid map of plasmid pPF14.

Figure 14 is a plasmid map of plasmid pCW186.

Figure 15 is a plasmid map of plasmid pCW187.

Figure 16 is a plasmid map of plasmid pPF66.

Figure 17 is a plasmid map of plasmid pPF57.

Figure 18 is a diagrammatic representation of an antisense gene construction for use in homeologous meiotic recombination.

Figure 19 is a plasmid map of plasmid p3243.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the inventors' discovery that there exist in higher plants genes which are homologous to MMR genes in *E. coli*, and to MMR genes in 30 yeasts and humans.

Thus, the inventors have identified genes, herein designated AtMSH3 and AtMSH6, of the plant Arabidopsis thaliana which encode the proteins AtMSH3 and AtMSH6. These plant proteins are homologous to yMSH3 and yMSH6, respectively. The present inventors have isolated cDNAs encoding the proteins AtMSH3 and AtMSH6 and have isolated the complete gene encoding AtMSH6. Given the teaching herein, other genes (for example AtMSH2, and genes of other plants) may be obtained which are involved in DNA mismatch repair in plants, including other genes which encode polypeptides homologous to MMR proteins of yeasts or humans, such as genes which encode

The state of the state of

Č

WO 99/19492 PCT/EP98/06977

polypeptides homologous to yeast MSH2. MLH1 or PMS2, or to human MLH1, PMS1 or PMS2. For example, given the teaching herein, genes of members of the *Brassicaceae* family or of other unrelated families, for example the *Poaceae*, the *Solanaceae*, the *Asteraceae*, the *Malvaceae*, the *Fabaceae*, the *Linaceae*, the *Canabinaceae*, the *Dauaceae* and the *Cucurbitaceae* family, and which encode polypeptides homologous to MMR proteins of yeasts or humans may be obtained.

Examples of plants whose genes encoding polypeptides homologous to MMR proteins of yeasts or humans may be obtained given the teaching herein include maize, wheat, oats, barley, rice, tomato, potato, tobacco, capsicum, sunflower, lettuce, artichoke, safflower, cotton, okra, beans of many kinds including soybean, peas, melon, squash, cucumber, oilseed rape, broccoli, cauliflower, cabbage, flax, hemp, hops and carrot.

Within the meaning of the present invention, a first polypeptide is defined as homologous to a second polypeptide if the amino acid sequence of the first polypeptide exhibits a similarity of at least 50% on the polypeptide level to the amino acid sequence of the second polypeptide.

A procedure which may be followed to obtain genes AtMSH3 and AtMSH6 is described in Example 1. Essentially the same technique may be applied to obtain other mismatch repair genes of Arabidopsis thaliana, and essentially the same technique as 20 exemplified herein may be applied to cDNA obtained by reverse transcription of RNA from other plants. Alternatively, given the sequence information disclosed herein, other degenerate oligonucleotide primers, especially oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions (such as the conditions described in Example 1 using primers UPMU and DOMU) to AtMSH3 and/or AtMSH6 25 may be designed and obtained for use in isolating sequences of plant mismatch repair genes which are homologous to AtMSH3 or AtMSH6, from other plants. oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions to plant mismatch repair genes of plants other than Arabidopsis thaliana also fall within the scope of the present invention and may be utilised to obtain mismatch 30 repair genes of still other plants. Typically, such oligonucleotides are capable of hybridising at 45°C under standard PCR conditions to a DNA molecule which encodes a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or a human. The temperature at which oligonucleotides of the invention hybridise to AtMSH3 and/or AtMSH6, or to plant mismatch repair genes of plants other than Arabidopsis thaliana, or 35 to DNA molecules which encode polypeptides which are homologous to a mismatch repair polypeptide of a yeast or a human may be higher than 45°C, for example at least 50°C, or at least 55°C, or at least 60°C or as high as 65°C.

The second secon

The successful gene isolation disclosed herein demonstrates for the first time the existence of MMR in higher plants and indicates the presence of other plant MMR genes. For example, genes encoding the plant homologs of MSH1, MSH2, MSH4, MSH5, PMS1, PMS2 and MLH1 may be identified given the teaching herein. Such genes, as well as those specifically described herein, separately or in combination, are useful in manipulating the plant MMR for plant breeding purposes. Thus, for example, the plant MMR may be altered by including in a plant cell a polynucleotide sequence as defined herein above with reference to the third embodiment of the invention, and causing the polynucleotide sequence to express either a polynucleotide which disables a plant MMR gene, or a polypeptide which disrupts the plant's MMR system.

The DNA molecule of the third embodiment of the invention includes a polynucleotide sequence (herein referred to as a MMR altering gene) which may for example encode sense, antisense or ribozyme molecules characterised by sufficient base sequence similarity or complementarity to the gene to be altered to permit the antisense or 15 ribozyme molecule to hybridise with the plant MMR gene in vivo or to permit the sense molecule to participate in co-suppression. Alternatively, the MMR altering gene may encode a protein or proteins which interfere with the activity of a plant MMR protein and thus disrupt the plant's MMR system. For example, such encoded proteins may be antibodies or other proteins capable of interfering with MMR protein function, such as by 20 complexing with a protein functionally involved in plant MMR thereby disrupting the MMR of the plant. An example of such a protein is the MSH3 protein of Arabidopsis thaliana described herein or a protein of another plant which is homologous to the MSH3 protein of A. thaliana. For instance, overexpression of MSH3 in a plant cell causes MSH2 present in the cell to be substantially completely complexed, disrupting the 25 mismatch repair mechanism or mechanisms in the cell which are functionally dependent on the presence of a complex of MSH2 with MSH6. Similarly, mismatch repair mechanisms which depend on the presence of a complex of MSH2 and MSH3 may be disrupted by the overexpression of MSH6.

A chimeric gene of the fourth embodiment, incorporating a MMR altering gene, 30 may be prepared by methods which are known in the art. Similarly, the MMR altering gene may be introduced into a plant cell, regenerating tissue or whole plant by techniques known in the art as being suitable for plant transformation, or by crossing. Known transformation techniques include Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters. Suitable promoters may direct constitutive expression, such as the 35S or the NOS promoter. Usually, however, the promoter will direct either inducible or tissue specific (e.g. callus; embryonic tissue; etc.), cell type specific (e.g. protoplasts; meiocytes; etc.) or developmental (e.g. embryo) expression of the altering gene or genes, in order for the

MMR system to function in tissue types or cell types, or at developmental stages of the plant, in which it is not desirable for the MMR system to be altered. Using such promoters, therefore, the activity of a MMR altering gene may be limited to a specific stage during plant development or it may be altered by controlling conditions external to the plant, and the deleterious effects of a permanently disabled or altered DNA mismatch repair system in a plant may be avoided. Examples of suitable promoters which are not constitutive are known in the art and include inducible promoters such as *PR*1a (reviewed by Gatz, 1997, Annual Rev. Plant Phys. Plant Mol. Biol. 48: 89), tissue specific promoters such as *AoPRI* (Sabahattin et al., 1993, Biotechnology 11: 218), and cell-type specific promoters such as *DMC*1.

A chimeric gene in accordance with the invention may further be physically linked to one or more selection markers such as genes which confer phenotypic traits such as herbicide resistance, antibiotic resistance or disease resistance, or which confer some other recognisable trait such as male sterility, male fertility, grain size, colour, growth rate, flowering time, ripening time, etc.

The process of the tenth embodiment of the invention provides, for example, a process for generating intraspecies genetic variation by altering the mismatch repair system in a plant cell, in regenerating plant tissue or in a whole plant. The plant cell, regenerating tissue or whole plant includes and expresses one or more MMR altering 20 genes which are capable of altering mismatch repair in the plant cell, regenerating tissue or whole plant. Alteration of MMR may be achieved, for example, by inactivating the genes encoding plant MSH3 and/or plant MSH6. It is preferred to inactivate the plant MSH3 and MSH6 encoding genes at the same time and in the same plant cell, regenerating tissue or whole plant. Typically in this preferred form of the invention 25 inactivation of either plant MSH3 or MSH6 alone is insufficient to substantially alter the plant's mismatch repair system and only when both MSH3 and MSH6 are inactivated simultaneously is the plant's mismatch repair system sufficiently altered to prevent the MMR system from recognising base pair mismatches, base insertions or deletions as a result of DNA replication errors, DNA damage, or oligonucleotide induced site-specific 30 mutagenesis. However, in some applications of the invention, inactivation of only one gene may also be used to cause genomic instability or increase the efficiency of sitespecific mutagenesis.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the genome of the plant cell, regenerating tissue or whole plant in order to restore mismatch repair in the plant cell, regenerating tissue or whole plant. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools, such as ribozymes, or may be removed from the genome using gene elimination systems known in the art, such as CRE/LOX. It is preferred to render two genes, whose gene products combine to incapacitate MMR, ineffective by separating

the contract of the contract

the altering genes through segregation. Therefore, in a preferred embodiment of the invention a first plant cell or plant is generated in which only plant MSH3 is incapacitated, and a second plant cell or plant is generated in which only plant MSH6 is incapacitated. The combination of both genomes, for example by crossing, then produces significant 5 MMR deficiency in those cells or plants which have inherited both altering genes. If the altering genes are expressed from unlinked loci, gene segregation restores MMR activity in the progeny of the cells or plants.

In a process of the ninth embodiment of this invention, homeologous recombination is enhanced between different genomes, chromosomes or genes in plant cells or plants by altering MMR in said plant cells or plants. Such genomes, chromosomes or genes are characterised in that they originate from different plant families, genera, species, subspecies, plant varieties or lines. Hybrid plant cells or hybrid plants may be produced by crossing, by cell fusion or by other techniques known in the art. These plant cells or plants are further characterised by expressing one or more genes that are capable of altering mismatch repair in the plant cell or plants.

In the process of the ninth embodiment, the homeologous recombination is typically for the purpose of introducing a desired characteristic in the hybrid plant. In this typical application of the process of the ninth embodiment, and in the process of the tenth embodiment the desired characteristic may be any characteristic which is of value to the plant breeder. Examples of such characteristics are well known in the art and include altered composition or quality of leaf or seed derived storage products (e.g. oil, starch, protein), altered composition or quality of cell walls (e.g. decrease in lignin content), altered growth rate, prolonged flowering, increased plant yield or grain yield, altered plant morphology, resistence to pathogens, tolerance to or improved performance under environmental stresses of various kinds, etc.

In a preferred form of the tenth embodiment, an MMR altering gene is cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant to produce a
hybrid plant cell or hybrid plant in which homeologous recombination can occur.

Suitably, the MMR proficient plant cell or MMR proficient plant may also include an
MMR altering gene. For example a gene capable of inactivating plant MSH3 may be cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant in which MSH6
is inactivated. A resultant hybrid plant in which homeologous recombination occurs will
include both the MSH3 and MSH6 altering genes and its MMR system will therefore be
inactivated.

In this form of the invention, if hybrid plants are to be produced by crossing, the MMR altering gene preferably originates from the male parent, thus ensuring that the

MMR altering gene is always introduced and is not present in the recipient cell. That is, the MMR of the recipient cell, prior to introduction of the MMR altering gene, is typically proficient. Alternatively, if an MMR altering gene is present in a recipient cell it may be ineffective or inefficient on its own, or it may be linked to an inducible or tissue specific or cell type specific promoter which only renders the MMR altering gene active under limited conditions.

Thus, in a preferred form of the process of the ninth embodiment, the MMR system of the hybrid plant is initially unaltered. In this form of the process, the step of altering the mismatch repair system may comprise introducing into the hybrid plant, or cells thereof, a MMR altering gene, such as by Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters, as described above. Preferably, the promoter is transcriptionally active in mitotically and meiotically active tissue and/or cells to ensure MMR alteration after chromosome pairing at mitosis and meiosis, respectively. The preferred timing for MMR alteration is at meiosis, because recombinant genomes, chromosomes or genes are directly transmitted to the progeny. A suitable meiocyte specific promoter is for example the *DMC*1 promoter from *Arabidopsis thaliana* ssp. *Ler.* (Klimyuk and Jones, 1997, Plant J. 11, 1-14). However, mitotic homeologous recombination is also a desirable outcome as somatic recombination events can be transmitted to offspring due to the totipotency of plant cells and the lack of predetermined germ cells in plants.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the hybrid plant or hybrid plant cells, in order to restore mismatch repair in the hybrid plant or hybrid plant cells. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools as described herein above.

EXAMPLES

Example 1. Cloning of the AtMSH3 and AtMSH6 coding sequences

Isolation of partial AtMSH3 and AtMSH6 consensus sequences

Degenerate oligonucleotides UPMU (SEQ ID NO:1) and DOMU (SEQ ID NO:2)

UPMU CTGGATCCACIGGICCIAA(C/T)ATG

DOMU CTGGATCC(A/G)TA(A/G)TGIGTI(A/G)C(A/G)AA

were used to isolate AtMSH3 and AtMSH6 sequences by PCR amplification.

Primers UPMU and DOMU correspond to conserved amino acid sequences of the proteins MutS (E. coli and S. typhimurium), HexA (S. pneumoniae). Repl (mouse) and Duc1 (human). The conserved regions to which they are targeted are TGPNM for UPMU (amino acid positions 852-856 for AtMSH6 and 816-820 for AtMSH3) FATHY or FVTHY

The second of th

for DOMU (amino acid positions 964-968 for AtMSH6 and 928-932 for AtMSH3, respectively.) These primers have been used to isolate MSH2 and MSH1 from yeast (Reenan and Kolodner, Genetics 132: 963-973 (1992)) and MSH2 from *Xenopus* and mouse (Varlet et al., Nuc. Acids Res. 22:5723-5728 (1994)).

Template single strand cDNA was produced by reverse transcription of 2 µg total RNA from a cell suspension culture of Arabidopsis thaliana ecotype Columbia (Axelos et al. 1989, Mol. Gen. Genetics 219: 106-112). The PCR reaction was performed under the following conditions in a final volume of 100µl: 0.2mM dNTP, 1µM each primer, 1XPCR buffer, Iu Tag DNA polymerase (Appligene) in the presence of template cDNA. PCR 10 parameters were 5 minutes at 94°C, followed by 30 cycles of 40 seconds at 95°C, 90 seconds at 45°C, I minute at 72°C. The amplification products were cloned into pGEM-T vector (Promega) and sequenced. Two different clones were isolated, S5 (350bp) was homologous to MSH3, S8 (327bp) was homologous to MSH6. Complete cDNA sequences were then isolated according to the Marathon cDNA amplification kit procedure (Clontech). 15 In summary, this procedure involves producing double stranded cDNA by reverse transcription of 2µg polyA+ RNA from the cell suspension culture of Arabidopsis. Adaptors are ligated on each side of the cDNA. The ligated cDNA is used as a template for 5' and 3' RACE PCR reactions in the presence of primers that are specific for the adaptor on one side (AP1 and AP2), and specific for the targeted gene on the other side. A 5' and a 3' 20 fragment that overlap are thus produced for each gene. The complete gene coding sequence can be reconstituted taking advantage of a unique restriction site. if available, in the overlapping region. Specific details of this procedure as it was used to isolate AtMSH3 and AtMSH6 coding regions, are as follows.

Isolation of AtMSH3 complete coding sequence

From the sequence of clone S5, primer 636 (SEQ ID NO:3) was designed:

636 TGCTAGTGCCTCTTGCAAGCTCAT.

Primer AP1 (SEQ ID NO:4) is complementary to a portion of an adaptor sequence which had been ligated to the 5' and 3' ends of *Arabidopsis* cDNA:

API CCATCCTAATACGACTCACTATAGGGC.

PCR performed on the ligated cDNA with primers 636 and AP1 for the 5' RACE PCR was followed by a second round of amplification with the nested primers AP2 (SEQ ID NO:5) and S525 (SEQ ID NO:6)

AP2 ACTCACTATAGGGCTCGAGCGGC

S525 AGGTTCTGATTATGTGTGACGCTTTACTTA

35 (the latter was also designed to correspond to a part of the sequence of clone S5) and produced a 2720bp DNA fragment. Figure 1 provides a diagrammatic representation of the primer sequences used to isolate *AtMSH3*. Another primer (S51, SEQ ID NO:7)

and the state of t

S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG

was designed closer to the 5° border and permitted the determination of 99bp upstream to the ATG initiation codon. For the 3° RACE PCR, a first PCR reaction was performed with primers AP1 and 635 (SEQ ID NO:8).

- 635 GCACGTGCTTGATGGTGTTTTCAC
- 5 followed by a second round of amplification, using the nested primers AP2 and S523 (SEQ ID NO:9)
- S523 TCAGACAGTATCCAGCATGGCAGAAGTA
 which produced a DNA fragment of 890bp. Both DNA fragments were subcloned into pGEM-T and sequenced. Since PCR amplification using the Expand Long Template PCR
 system (Boehringer-Mannheim) produced errors in the sequence, new oligonucleotides were designed to isolate those sequences again by PCR, but with the high fidelity DNA polymerase Pfu. PCR with primers 1S5 (SEQ ID NO:10) and S53 (SEQ ID NO:11)
 - 1S5 ATCCCGGGATGGGCAAGCAAAAGCAGCAGACGA
 - S53 GACAAAGAGCGAAATGAGGCCCCTTGG
- amplified the 1244bp fragment clone 52 (SEQ ID NO:12, cloned into pUC18/Sma1). PCR with primers S52 (SEQ ID NO:13) and 2S5 (SEQ ID NO:14)
 - 2S5 ATCCCGGGTCAAAATGAACAAGTTGGTTTTAGTC
 - S52 GCCACATCTGACTGTTCAAGCCCTCGC
- amplified the 2104bp clone 13 (SEQ ID NO:15, cloned into pUC18/Sma1). The complete coding sequence of the AtMSH3 gene was reconstructed in pUC18 by ligating the 5' half of AtMSH3 (clone 52) to the 3' half of AtMSH3 (clone 13) after digesting with BamH1 which has a unique cleavage site in the overlapping region of both clones. This manipulation yielded plasmid pPF26. The SmaI fragment from pPF26 contains the complete AtMSH3 coding sequence. The remaining primers referred to in Figure 1 are as follows:
 - S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG (SEQ ID NO:16)
 - S525 AGGTTCTGATTATGTGTGACGCTTTACTTA (SEQ ID NO:17)

Figures 2 and 3 provide plasmid maps of clones 52 and 13 respectively, showing restriction enzyme cleavage sites. The complete *AtMSH3* coding sequence (SEQ ID NO:18) 30 is 3246bp long and is shown in Figure 4 together with the deduced sequence (SEQ ID NO:19) of the encoded polypeptide. *AtMSH3* is clearly homologous to the yeast and mouse *MSH3* genes. A sequence alignment of polypeptides encoded by *AtMSH3* and that encoded by *Saccharomyces cerevisiae MSH3* is set out in Figure 5.

Isolation of the AtMSH6 complete coding sequence and genomic sequences

The same procedure allowed isolation of the *AtMSH6* cDNA. Figure 6 provides a diagrammatic representation of the primer sequences used to isolate *AtMSH6*. For the 5' RACE PCR, primers 638 (SEQ ID NO:20) and AP1 (SEQ ID NO:4)

638 TCTCTACCAGGTGACGAAAAACCG allowed the amplification of a 2889 DNA fragment. Primer S81 (SEQ ID NO:21)

WO 99/19492 PCT/EP98/06977

S81 CGTCGCCTTTAGCATCCCCTTCCTTCAC

helped define the 142bp upstream to the ATG initiation codon. On the 3' side, RACE PCR was initially performed with primers S823 (SEQ ID NO:22) and AP1 (SEQ ID NO:4),

S823 GCTTGGCGCATCTAATAGAATCATGACAGG
5 and then with the nested primers 637 (SEQ ID NO:23) and AP2 (SEQ ID NO:5).

637 GACAGCGTCAGTTCTTCAGAATGC

to produce a 774bp DNA fragment. As for *AtMSH3*, those fragments were cloned and sequenced. Re-isolation of the DNA sequence using the high fidelity *Pfu* polymerase and newly designed primers 1S8 (SEQ ID NO:24) and S83 (SEQ ID NO:25) (for the 5' side) led to a 2182 bp DNA fragment identified as clone 43 (SEQ ID NO:26, cloned in pUC18/Sma1), and a 1379bp clone identified as clone 62 (SEQ ID NO:27, also cloned in pUC18/Sma1).

	1S8	ATCCCGGGATGCAGCGCCAGAGATCGATTTTGT
	2S8	ATCCCGGGTTATTTGGGAACACAGTAAGAGGATT (SEQ ID
i	NO:28)	
	S82	GCGTTCGATCATCAGCCTCTGTGTTGC (SEQ ID NO:29)
	S83	CGCTATCTATGGCTGCTTCGAATTGAG

Figures 7 and 8 provide plasmid maps of clones 43 and 62 respectively, showing restriction enzyme cleavage sites. Clones 43 and 62 were digested by the *Xmn*1 restriction enzyme for which a unique site is present in their overlapping region and then ligated. The complete *AtMSH6* coding sequence (SEQ ID NO:30) is 3330bp long and is shown in Figure 9 together with the deduced sequence (SEQ ID NO:31) of the encoded polypeptide. *AtMSH6* is clearly homologous to the yeast and mouse *MSH6*genes. A sequence alignment of polypeptides encoded by *AtMSH6* and that encoded by *Saccharomyces cerevisiae MSH6* is set out in Figure 10.

An AtMSH6 genomic sequence was also isolated from a genomic DNA library constituted after partial Sau3AI digestion of DNA from the Arabidopsis cell suspension. 8062bp were sequenced that covered the AtMSH6 gene and show colinearity with the cDNA. 16 introns are found scattered along the gene. The complete genomic sequence 30 (SEQ ID NO:98) is shown in Figure 11.

Example 2. A measure of somatic variation in MMR deficient plants Constructs

Constructs with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 under the control of a single 35S promoter have been inserted into the binary vector pPZP121 (Hajdukiewicz et al., Plant Mol. Biol. 23, 793-799) between the right and left borders of the T-DNA. The pPZP121 plasmid confers chloramphenicol resistance to Escherichia coli or Agrobacterium tumefaciens bacteria. The aacC1 gene is carried by the T-DNA and allows selection of transformed plant cells on gentamycin (Hajdukiewicz et al., Plant Mol. Biol. 25, 989-994). For the purpose of expressing antisense constructs, a 35S

Part Rose The trees The state of ij. The state of the s

promoter/terminator cassette with a polylinker was introduced into pPZP121. The 3' ends of the considered genes have been chosen since this region seems more efficient for antisense For AtMSH3 this corresponds to clone 13 (2104bp), for AtMSH6 this corresponds to clone 62 (1379bp). Clone 13 comprises 2104bp of the 3' region that were cut 5 off the pUC18 vector by Sal1/Sst1 restriction, blunted with T4 DNA polymerase and ligated into the T4 DNA polymerase blunted BamHI site of pPZP121/35S, creating clone pPF13. The same procedure was followed for the 3' region of AtMSH6 clone 62 (1379bp) thus creating plasmid pPF14. For the double constructs, the 3' region (from clone 62) of AtMSH6 was introduced ahead of the AtMSH3 region into pPF13 creating pCW186 and 10 reciprocally, the 3' region of AtMSH3 (from clone 13) was introduced ahead of AtMSH6 into pPF14, creating pCW187.

These constructs were introduced into the Arabidopsis cells (as described below) of wildtype Columbia and of the Columbia tester line.

An alternative strategy to antisense inhibition of AtMSH6 comes from experiments of 15 Marra et al. (1998. Proc. Natl. Acad. Sci USA 95, 8568-8573) who show that overexpression of functional MSH3 results in depletion of MSH6 protein in human cells. This depletion may generate a mismatch repair mutant phenotype.

For the purpose of overexpressing functional AtMSH3 protein in plant cells, the complete MSH3 coding region was excised from pPF26 (example 1) by digestion with 20 Smal, and was inserted into the Smal site of pCW164. The resulting construct was named pPF66. It contains a complete AtMSH3 gene under the control of the 35S promoter inside the left (LB) and right (RB) border of the T-DNA. This T-DNA also contains the hpt2 gene for gentamycin selection. Plasmid pPF66 was introduced into Arabidopsis cells as described below. One cell clone was selected which clearly overexpressed the AtMSH3 25 gene as shown by Northern analysis. Figures 12-16 provide plasmid maps of plasmids pPF13, pPF14, pCW186, pCW187 and pPF66, respectively.

Construction of tester construct

For the purpose of Forward Mutagenesis Assays, a tester construct was built containing the coding regions for nptII, codA, uidA. All three genes are driven by the 35S 30 promoter and are terminated by the 35S terminator. This construct was obtained by introducing an EcoR1 fragment encoding the codA cassette (2.5kb) and a HindIII fragment encoding the uidA (GUS) cassette (2.4kb) into the pPZP111 vector (Hajdukiewicz et al.,1994, Plant Mol Biol 23: 793-799) which already contained the nptII expression cassette. This new plasmid was named pPF57. NptII is used to select for transformed plant cells. 35 GUS is used to analyse the degree of gene silencing in the construct (i.e. to identify cell lines in which the transgenes are expressed), and codA is used as a marker for forward mutagenesis (described below).

The state of the s

The plasmid map of pPF57 is provided in Figure 17.

Plant cell transformation

The constructs are introduced into Agrobacterium by electroporation. Plant cells are then transformed by co-cultivation. A suspension culture of Arabidopsis thaliana cells that 5 has been established by Axelos et al. (1992, Plant Physiol. Biochem. 30, 1-6) may be used. One day old freshly subcultured cells are diluted five times into AT medium (Gamborg B5 medium. 30g/l sucrose, 200µg/l NAA). 10µl of saturated Agrobacterium containing the transforming T-DNA constructs are added to 10ml diluted cells in a 100ml erlenmeyer. The co-cultivation is agitated slowly (80rpm) for 2 days. The cells are then washed 3 times into AT medium and finally resuspended in the same initial volume (10ml). The culture is agitated for 3 days to allow expression before plating on selection plates (AT/BactoAgar 8g/l+gentamycin 50µg/ml). Transformed individual calli are isolated 3 weeks later.

Tester Strain

The tester construct on plasmid pPF57 was introduced into Arabidopsis cells of wildtype Columbia using the transformation protocol described above. Among 10 candidate transformants, one cell clone was shown (by Southern analysis) to have a unique T-DNA insertion. All three genes were shown to be functional in this cell line as indicated by resistance to kanamycin, blue staining in the presence of X-Glu (GUS), and sensitivity to 5-fluoro-cytosine (codA).

MMR altering genes (described above) were then introduced individually into the tester line and transformed cells are used for analysis of both Microsatellite Instability and Forward Mutagenesis.

Microsatellite analysis

Microsatellites have been described in Arabidopsis (Bell and Ecker, 1994, Genomics 19, 137-144). The present Example is based on a study of instability of microsatellites that are polymorphic amongst different ecotypes. DNA is extracted from the transformed calli. Specific primers have been defined that are used to amplify the microsatellite sequence. One of the two primers is previously P³² labelled by T4 kinase. In case of a polymorphic variation, new PCR products appear that do not follow the expected pattern of migration on a polyacrylamide gel. This is a commonly observed feature for MMR deficient cells in yeast or mammalian cells.

In particular, the present Example describes a study on microsatellites ca72 (CA₁₈), nga172 (GA₂₉), and ATHGENEA(A₃₉), chosen because they belong to the types predominantly affected in human mismatch repair deficient tumors. The size of these microsatellites is not conserved from one *Arabidopsis* ecotype to the other.

Arabidopsis cells which are transformed with an MMR altering gene (above) and control cells not expressing the MMR altering gene are allowed to form calli. DNA is

the type of the contract of th

1 1 1

WO 99/19492

10

rapidly extracted from the calli and is analysed for microsatellite instability as described in detail by Bell and Ecker 1994. Genomics 19, 137-144. In summary, the relevant microsatellite is amplified by PCR using P32 labelled primers. The PCR products are separated on a DNA sequencing gel for size determination. Size differences between microsatellites from transformed and control cells not expressing the MMR altering gene in question indicate microsatellite instability as a result of MMR alteration.

The sequences of primers used for PCR amplification of microsatellites ca72 and nga172 are included in Table 1. PCR amplification of microsatellite ATHGENEA made use of a forward primer containing the sequence

ACCATGCATAGCTTAAACTTCTTG (SEQ ID NO:32)

and of a reverse primer containing the sequence

ACATAACCACAAATAGGGGTGC (SEQ ID NO:33).

The amplification for microsatellite ca72 revealed in Columbia control cells (with respect to the MMR altering gene) a 248 bp long PCR fragment instead of the published length of 124 bp. DNA sequencing verified this fragment as a CA₁₈ microsatellite.

Forward mutagenesis assay

Tester cells transformed with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 are analysed for the stability of the codA gene. The functional codA gene confers to sensitivity to 5-fluoro-cytosine (5FC), whereas a gene inactivating mutation in codA will confer resistance to 5FC. The frequency of resistant cells is therefore a good indicator of somatic variation as a direct result of MMR alteration. Variants resistant to 5FC are first analysed for GUS activity. If GUS is mactive, 5FC resistance is assumed to be due to gene silencing (all three genes are under the 35S promoter). If GUS is active, 5FC resistance is assumed to be due to forward mutations that have inactivated codA. PCR is then performed on the putative codA mutant genes which is then sequenced to confirm the presence of forward mutations in codA.

Besides *codA*, other marker genes may also be used for the Forward Mutagenesis Assay such as the *ALS* gene (conferring sensitivity to valine or to sulfonylurea; Hervieu and Vaucheret, 1996, Mol. Gen. Genet. 251 220-224; Mazur et al. 1987, Plant Physiol. 85 1110-30 1117).

Example 3. Homeologous meiotic recombination in Arabidopsis thaliana

- A. Construction of a meiocyte specific gene expression cassette comprising the *DMC*1 promoter and the *NOS* terminator
- (i) The *DMC*1 promoter may be used as published by Klimyuk and Jones, 1997, 35 Plant J. 11,1-14). To obtain a more convenient alternative for gene cloning, a 3.3 Kb

long subfragment of the *DMC*1 promoter was obtained by PCR from genomic DNA of *Arabidopsis thaliana* (ssp. Landsberg erecta "Ler").

The PCR was done in three rounds:

Round One: A 3.7 Kb long product was obtained using the forward primer 5 DMCIN-A comprising the sequence

GAAGCGATATTGTTCGTG (SEQ ID NO:34)

and the reverse primer DMCIN-B comprising the sequence

AGATTGCGAGAACATTCC (SEQ ID NO:35).

The weak amplification product was then used as template for round two and three.

Round Two: A 3.1 Kb long product comprising the promoter and the 5' untranslated leader was obtained using forward primer DMCIN-1, which contained the sequence

acgcgtcgacTCAGCTATGAGATTACTCGTG (SEQ ID NO:36)

and introduced a SaII cloning site at the 5' end of the promoter fragment, and reverse primer DMCIN-2 which contained the sequence

getetagaTTTCTCGCTCTAAGACTCTCT (SEQ ID NO:37)

and introduced a XbaI site at the 3' end of the PCR fragment.

Round Three: A 0.2 Kb long product comprising the first exon/intron of the *DMC*1 promoter was obtained using forward primer DMCIN-3, which contained the sequence

getetagaGCTTCTCTTAAGTAAGTGATTGAT (SEQ ID NO:38)

and introduced a XbaI site at the 5' end of the PCR fragment, and reverse primer DMCIN-4, containing the sequence

tecceegggetegagagatetecatggTTTCTTCAGCTCTATGAATCC (SEQ ID NO:39) and introduced at the 3' end of the PCR product restriction sites for Ncol, Bg/II. Xhol and 25 Smal.

The products obtained in round Two and Three were digested with XbaI and subsequently ligated to reconstitute a 3.3 Kb long DMC1 promoter from which the first two in-frame ATG start codons were replaced with a unique restriction site for XbaI. This promoter can be cloned between the restriction sites for SaII and SmaI of p3264, which contains the SacI-EcoRI NOS terminator in pBIN19, to yield the entire expression cassette in pBIN19. This cassette contains the following cloning sites: NcoI, BgIII, XhoI. SmaI and (already present on p3264) KpnI and SacI.

(ii) Another strategy yielded the following convenient *DMC*1 promoter. A 1.8 kb DNA fragment comprising the 3' terminal part of the meiocyte specific *DMC*1 promoter was isolated by PCR from purified genomic DNA of *Arabidopsis thaliana* (ssp. Landsberg erecta "*Ler*"). The forward PCR primer (DMC1a) contained the sequence

acgcgtcgacGAATTCGCAAGTGGGG (SEQ ID NO:40)

and introduced a Sall cloning site at the 5' end of the promoter fragment. The reverse PCR primer (DMC1b) contained the sequence

the same of

The second of th

tecatggagatetecegggtacCGATTTGCTTCGAGGG (SEQ ID NO:41)

introducing a polylinker region at the 3' end of the promoter fragment. The PCR reaction was carried out with VENT DNA Polymerase (NEB) over 25 cycles using the following cycling protocol: 1 minute at 94°C, 1 minute at 54°C, 2 minutes at 72°C.

The PCR reaction yielded a blunt ended DNA fragment which was digested with restriction enzyme SalI and was cloned into the cleavage sites of restriction enzymes SalI and SmaI in plasmid p2030, a pUC118 derivative containing the SacI-EcoRI NOS terminator fragment of pBIN121. The cloning yielded plasmid p2031, containing the DMC1 promoter-polylinker-NOS terminator expression cassette depicted in Figure 18.

to B. Construction of an MSH3 antisense gene under the control of the DMC1 promoter

A 2.1 kb DNA fragment encoding the carboxyterminal part of AtMSH3 was removed from the polylinker of clone 13 described in Example 1 after (i) digestion with *Kpn*I. (ii) blunting of the DNA ends generated by *Kpn*I and (iii) digestion with *BamHI*. The isolated fragment was then cloned in antisense orientation downstream of the *DMC*1 promoter in plasmid p2031, which had been digested with restriction enzymes *SmaI* and *BgIII*. This cloning yielded plasmid p2033 (Figure 18).

After digestion of p2033 with *EcoRI*. a 4.1 kb DNA fragment was recovered comprising the *DMC1* promoter, the partial *MSH3* cDNA in antisense orientation with respect to the promoter and the *NOS* terminator. This fragment was cloned into the *EcoRI* restriction site of plant transformation vector pNOS-Hyg-SCV to yield plasmid p3242 (Figure 18).

C. Construction of a combined MSH6/MSH3 antisense gene under the control of a single DMC1 promoter

A 3.1 kb fragment, encoding in antisense orientation the partial AtMSH6 and AtMSH3 sequences and the 35S terminator, was isolated from pCW186 by digestion with KpnI. The fragment was treated with Klenow enzyme to blunt both ends. It was then cloned into the SmaI site of plasmid p3243 (a pNOS-Hyg-SCV derivative, illustrated in Figure 19), which had been opened to delete the region between the SmaI sites. Clones containing the fragment in the antisense orientation with respect to the DMC1 promoter (described in A(ii) above) were identified by diagnostic digestion with BamHI. The selected construct was named p3261.

Another practical way of cloning the double antisense gene is as follows. A 1 kb DNA fragment encoding the carboxyterminal part of AtMSH6 is isolated from clone 62 described in Example 1 after digestion of clone 62 plasmid DNA with BamHI, which cleaves in the 5' polylinker region flanking the partial cDNA, and with EcoRI, which cleaves within the cDNA. The isolated fragment is treated with Klenow enzyme to blunt both its ends and is cloned into the recipient plasmid p2033 or p3242. For the purpose of

The state of the state of

cloning, the recipient plasmid may be cleaved with either AvaI or NcoI and can be blunted with Klenow enzyme to produce blunt acceptor ends for fragment cloning. This cloning yields two opposite orientations of cloned fragment DNA with respect to the DMC1 promoter. These can be identified by diagnostic digestion with restriction enzymes ScaI or XmnI in conjunction with SacI. The selected construct contains the DMC1 promoter, the combined partial cDNAs for AtMSH3 and AtMSH6 (both cloned in antisense orientation with respect to the DMC1 promoter) and the NOS terminator. If the recipient plasmid is p2033, the combined antisense gene under control the single DMC1 promoter is recovered from the vector after EcoRI digestion and is cloned into the EcoRI restriction site of pNOS-Hyg-SCV.

D. Construction of a full-length MSH3 sense gene under control of the DMC1 promoter for overexpression of functional MSH3 protein

Overexpression of MSH3 protein was shown in human cells (Marra et al., 1998, Proc. Natl. Acad. Sci. USA 95, 8568-8573) to complex all available MSH2 protein. This leaves MSH6 protein without its partner, leading to the degradation of MSH6 protein and eventually to a mismatch repair phenotype.

This phenomenon is exploited to increase homeologous meiotic recombination in Arabidopsis as an alternative to antisense inhibition of *AtMSH*6. For this purpose the full-length cDNA encoding *AtMSH*3 is isolated from plasmid pPF66 by digestion with *SmaI* and is cloned into the *SmaI* site of the *DMC*1 expression cassettes described in A(i).

E. Selection of Recombination markers on homeologous chromosomes of *Arabidopsis* thaliana subspecies Landsberg erecta (Ler), Columbia (Col) and C24, respectively

E(i). Visual recombination markers in Arabidopsis th. subspecies C24:

Agrobacterium mediated transformation with a T-DNA containing a 35S-GUS gene, inactivated by insertion of a 35S-Ac transposable element (Finnegan et al., 1993, Plant Mol. Biol. 22, 625-633), had yielded a C24 line in which the T-DNA construct was integrated into chromosome 2. Genetic and molecular analysis of this line shows that the Ac transposon had excised from its T-DNA locus thereby restoring GUS activity, but had re-inserted into the chromosome at a distance of 16.4 cM, where it stayed fixed (due to disablement of Ac) within the chlorina gene. Insertional inactivation of the chlorina gene caused a bleached phenotype in those plants that were homozygous for this mutation. Because of the two linked phenotypic markers, chlorina3A:Ac and GUS, this C24 line was used in crosses to wildtype Ler for analysis of meiotic homeologous recombination on chromosome 2 in conjunction with molecular recombination markers.

35 E(ii). Visual recombination markers in Arabidopsis th. Ler:

The Ler line NW1 (obtained from NASC, Nottingham, UK) contains one recessive visual marker per chromosome. i.e. an-1 on Chr.1, py-1 on Chr.2, gl1-1 on Chr.3, cer2-1

on Chr.4, and *ms1-1* on Chr.5. This line is used in crosses to wildtype *C24* which expresses an MMR altering gene for analysis of meiotic homeologous recombination on chromosomes 1-5 in conjunction with molecular recombination markers listed in Table 1.

Other Ler lines from NASC have several visual markers in close proximity to each other on the same chromosome. When these lines are used for hybrid production, analysis of homeologous meiotic recombination may make use entirely of visual recombination markers. Particularly suitable for crossing to C24 wildtype that is expressing a MMR altering gene are the following Ler lines:

NW22: relative markers are dis1 - (4 cM) - ga4 - (11 cM) - th1 on chromosome 1.

NW10: relevant markers are tz-201 - (5 cM) - cer3 on chromosome 5.

NW134, relevant markers are ttg - (4 cM) - ga3 on chromosome 5.

NW24 (abi3-1) and NW64 (gl1-1). When present in the same plant on chromosome 3, abi3-1 and gl1-1 are 13 cM apart. Since this marker combination is not available from NASC, we have combined these markers by crossing line NW24 to line NW64. The F1 offspring were allowed to self-fertilise and to produce F2 seeds. F2 Plants which carry both markers as homozygous traits on the same chromosome can be identified firstly, by germinating F2 seeds on germination medium containing selective concentrations of abscisic acid, and subsequently, by identifying among the abscisic acid resistant plants those individuals which show the glabra phenotype.

20 E(iii) Molecular recombination markers in Col, Ler and C24:

The genome of Arabidopsis thaliana is interspersed with unique base sequences arranged as simple tandem repeats. Allelic repeats can vary in length between different Arabidopsis subspecies and when amplified by PCR yield diagnostic DNA products of different length named Simple Sequence Length Polymorphisms (SSLPs). Many SSLPs have been genetically mapped and have been assigned to unique chromosome locations on the recombinant inbred map (Bell and Ecker, 1994, Genomics 19, 137-144; Lister and Deans lines, Weeds World 4i, May 1997).

In Table 1 are listed 28 mapped and established SSLPs between *Ler* and *Col*. A number of PCR primer pairs are described herein (SEQ ID NO:42 to SEQ ID NO:97) 30 which also yielded SSLPs between *C24* and *Ler* (19 SSLPs) or between *C24* and *Col* (25 SSLPs), respectively. Polymorphic SSLPs can be used as molecular markers in the analysis of homeologous recombination between genomes from these subspecies.

The PCR reactions (25 µL) were carried out over 35 cycles (15 seconds at 94°C, 30 seconds at 55°C and 30 seconds at 72°C), with 0.25 U Taq DNA polymerase and 0.6 µg 35 genomic DNA in reaction buffer containing 2 mM MgCl₂. PCR products were separated by agarose gel electrophoresis (4% ultra high resolution agarose) and visualised by ethidiumbromide staining. The results from the PCR experiments are summarised in

Table 1, which also shows the sequence of PCR primers, primer annealing temperature (Tm), PCR product length and chromosome location of SSLP (with respect to the RI map of May 1997, Weeds World 4i).

F. Production of hybrid plants

C24 plants heterozygous for chlorina3A:Ac/GUS are crossed as male to emasculated wildtype Ler to produce Ler/C24(chlorina3A, GUS) hybrid seeds.

Due to the heterozygosity of the C24 parent, only 50 % of hybrid plants have inherited the chlorina3A:Ac/GUS locus. The remaining 50% of hybrid plants are wildtype with respect to chlorina3A:Ac/GUS. Since the mutant locus is linked to a kanamycin resistance gene (contained on the same T-DNA as GUS) mutant plants can be pre-selected by germinating hybrid seeds on germination medium containing 50 mg/L kanamycin.

Let plants homozygous for the five chromosome markers are male sterile (ms1-1) and are crossed without emasculation to wildtype C24 to produce Ler(an-1, py-1, gl1-1, cer2-1, ms1-1)/C24 hybrid seeds.

Other Ler plants, which are male fertile, are crossed after emasculation of the female parent to produce Ler/C24 hybrid seeds.

G. Introduction of MSH3 and MSH6/3 antisense genes into Arabidopsis and analysis of meiotic homeologous recombination

(i) Transformation of hybrid plants and analysis of homeologous meiotic recombination

The plant transformation vectors comprising the antisense genes described in (B) and (C) above are introduced into Agrobacterium tumefaciens strain AGL1 (Lazo et al., 1991, Bio/Technology 9, 963-967) by electroporation. Recombinant Agrobacterium clones are selected on LB medium containing 50 mg/L rifampicin and 100 mg/L carbenicillin. Selected clones are used to infect roots of Arabidopsis hybrid plants (described in (F) above) using the root transformation protocol of Valvekens et al. (1988, PNAS 85, 5536-5540) except that shoot and root inducing media contain hygromycin (10 mg/L) instead of kanamycin.

Plants regenerated from roots of hybrid plants are genetic clones of root donating plants and therefore are again genetic hybrids of two *Arabidopsis* subspecies described in 30 (F). However, in contrast to the root donating plants, the regenerated hybrid plants also contain the introduced transgene and the co-introduced hygromycin resistance gene with the latter allowing these plants to grow on hygromycin containing culture medium.

Hygromycin resistant plants are then allowed to enter the reproductive phase and to produce gametes by meiotic divisions of microspore and megaspore mothercells. At meiosis, the *DMC*1 promoter is activated and can direct the expression of antisense genes described in (B) and (C) above, leading to decreased *MSH*6 and/or *MSH*3 gene

WO 99/19492

expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3 protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between homeologous chromosomes.

Transgenic plants are then allowed to self-fertilise and to produce seeds. These 5 seeds (F2 seeds with respect to hybrid production), and the plants derived therefrom, carry the homeologous recombination events which can be identified by using the visual and molecular recombination markers described in (E) above.

In case of homeologous recombination between chromosomes of *Ler* and *C24(chlorina3A:Ac, GUS)*, the analysis concentrates on chromosome 2 by selecting plants showing the visual phenotypic marker *chlorina*. This marker thus serves as a reference point as it indicates that respective chromosomes 2 originate from *C24*. Other markers, such as *GUS* or molecular markers, on chromosome 2 may then be used to identify chromosomal regions which are derived from the *Ler* chromosome as a result of homeologous recombination. F2 plants of control transformants not expressing the antisense gene(s) can be analysed in parallel and the results can be used for comparison to homeologous recombination results obtained in antisense plants.

(ii) Transformation of C24 wildtype, hybrid plant production and analysis of homeologous meiotic recombination

Introduction of MMR altering genes into wildtype C24 is done using the root transformation protocol as described in G(i) for transformation of hybrid plants. Transformed plants are selected by resistance to either 10 mg/L hygromycin (in case of transformation with T-DNA's derived from pNOS-Hyg-SCV) or to 50 mg/L kanamycin (in case of transformation with T-DNA's derived from pBIN19).

Transgenic plants are then allowed to self-fertilise and to produce seeds (T1 seeds).

Segregation of the antibiotic resistance gene in the T1 population then indicates the number of transgene loci. Lines with a single transgene locus (indicated by a 3:1 ratio of resistant:sensitive plants) are selected and are bred to homozygosity. This is done by collecting selfed seeds (T2) from T1 plants and subsequent testing of at least four independent T2 seed populations for segregation of the antibiotic resistance gene. T2 populations which do not segregate the antibiotic resistance gene are assumed to be homozygous for both the resistance gene and the linked MMR altering gene.

C24 plants homozygous for the MMR altering gene are then crossed to Ler lines homozygous for recessive visual markers (see E(ii)) to produce C24/Ler hybrid plants as described in (F). These F1 hybrids are then allowed to enter the reproductive phase and to produce gametes by meiotic division of microspore and megaspore mothercells. At meiosis, the DMC 1 promoter is activated and can direct the expression of antisense or sense genes described in (B), (C) and (D) above, leading to decreased MSH6 and/or MSH3 gene expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3

protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between the homeologous chromosomes of C24 and Ler. Recombination events are then scored in the F2 generation.

For recombination analysis, the hybrid plants are allowed to self-fertilise and to produce F2 seeds. F2 plants are then preselected for a first visual marker. Since this marker is recessive, its visual presence indicates homozygosity for Ler DNA at the relevant locus. Those F2 plants which show this first visual marker are then analysed for the presence or absence of a second visual marker which in the Ler parent is closely linked to the first marker. Absence of the second visual marker indicates recombination between the relevant C24 and Ler chromosomes between the first and second marker. The frequency of recombination in transgenic hybrids is compared to the recombination frequency in control hybrids not expressing the MMR altering gene.

Examples of recombination analysis are the following.

The Ler line NW22(dis1, ga4, th1) is used for crosses to transformedC24.

F2 plants are preselected first for thiamine requirement (thI) and then are further analysed for re-appearance of wildtype height (loss of ga4) and/or re-appearance of normal trichomes (loss of disI) as a result of recombination.

The Ler line NW10(tz-201, cer3) is used for crosses to transformed C24.

F2 plants are then preselected first for thiazole requirement (tz) and then are further 20 analysed for re-appearance of normal, i.e. non-shiny stems (loss of cer3) as a result of recombination.

The Ler line NW134 (ttg, ga3) is used for crosses to transformed C24. F2 plants are first preselected for dwarfish appearance (ga3) and are then analysed for re-appearance of trichomes (loss of ttg) as a result of recombination.

Ler plants homozygous for abi3-1 and gl1-1 are used for crosses to transformed C24. F2 plants are first preselected for their ability to germinate in the presence of abscisic acid and are then analysed for re-appearance of trichomes on the leaves (loss of gl1-1) as a result of recombination.

In the case of homeologous recombination between transformed C24 and the Ler line 30 NW1 (an-1, py-1, gl1-1, cer2-1, ms1-1), recombination analysis is similar the one described above, except that the second marker is not a visual marker but has to be a molecular marker. This is because the Ler parent carries only one visual marker per chromosome.

receipting the receipt of the re-

Ę		
		Ì
	į	Will Street
		1
		Ž
E,	900	į
	7:22	1
	í	j
Ε		
	:::	=
	-	7
£		Į.
÷.,	-	illian i
Shade Shad	;=; ;=;	1

	L	TABLE 1: SS	BLE 1: SSLP Markers in Arabidopsis thaliana Subspecies	aliana S	ubspecies		
Chromosome	- (PCR Primer	Primer Sequence	Tm [°C]	length/COL	length/LER	length/C24
	Position	ran					
-	2.3	ATEAT! F	GCCACTGCGTGAATGATATG	57.8	172	162	162
	62	1 -	CGAACAGCCAACATTAATTCCC	58.2			
	9.3	NGA63 F	AACCAAGGCACAGAAGCG	57.3	111	68	120
-		NGA63 R	ACCCAAGTGATCGCCACC	59 6			
-	40.1	NGA248 F	TACCGAACCAAAACACAAAGG	56.1	143	129	no amplitic.
		NGA248 R	TCTGTATCTCGGTGAATTCTCC	58.2			
-	81.2	NGA128 F	GGTCTGTTGATGTCGTAAGTCG	60.1	180	061	no amplific.
	7:10	NGA128 R	ATCTTGAAACCTTTAGGGAGGG	58.2			
-	01.3	NGA280 F	CTGATCTCACGGACAATAGTGC	60.1	105	85	85
	91:5	NGA280 R	GGCTCCATAAAAAGTGCACC	57.8			
-	111 4	NGA111 F	CTCCAGTTGGAAGCTAAAGGG	09	128	162	170
-		NGA111 R	TGTTTTTAGGACAAATGGCG	70			
	ca 7.5	NGA 168 F	CCTICACATCCAAAACCCAC	57.8	213	217	208
		NGA 168 R	GCACATACCCACAACCAGAA	57.8			
				İ			

The second secon

Equ.
ŧ.
f.
1
Fig
E
1
T.J
*: E
£.j
C.

II ca							
	ca. 48	NGA1126L	CGCTACGCTTTTCGGTAAAG	57.8	161	661	961
		NGA1126R	GCACAGTCCAAGTCACAACC	59.9			
11 62	62.2	NGA361L	AAAGAGATGAGAATTTGGAC	51.7	114	120	114
		NGA361R	ACATATCAATATATAAAGTAGC	49.5			
11 73		NGA168 F	TCGTCTACTGCACTGCCG	9.69	151	135	135
		NGA168 R	GAGGACATGTATAGGAGCCTCG	61.9			
II Ca.	.77	AthBIO2 L	TGACCTCCTCTTCCATGGAG	. 6 65	141	209	139
		AthBIO2 R	TTAACAGAACCCAAAGCTTTC	54.5			
II	ca. 83	AthUBIQUE L	AGGCAAATGTCCATTTCATTG	54.1	146	148	148
		AthUBIQUE R	ACGACATGCAGATTTCTCC	57.8			
3.4	4	NGA172 F	AGCTGCTTCCTTATAGCGTCC	09	162	136	140
		NGA172 R	CATCCGAATGCCATTGTTC	55.4			
111 12	12.8	NGA126 F	GAAAAACGCTACTTTCGTGG	56.1	119	147	no amplific.
		NGA126 R	CAAGAGCAATATCAAGAGCAGC	58.2			
71 117	17.5	NGA162 F	CATGCAATTTGCATCTGAGG	55.8	107	68	по атрійс.
		NGA162 R	CTCTGTCACTCTTTCCTCTGG	1.09			

vij
L.
Fij
II.
ii.
E !
h-L
ri.
Fa., []

	0 10	NGA6 F	TGGATTTCTTCCTCTTCAC	56.1	143	123	143
	01.0	NGA6 R	ATGGAGGCTTACACTGATC	56.1			
		N OVON					
			OLOCIO OLOCATOR Oloca	6 65	247	234	220
N.	8.61	NGA12 F	AAIGIIGICICCCICCIC				
		NGA12 R	TGATGCTCTCTGAAACAAGAGC	58.2			
17/	24.1	NGA8 F	GAGGCCAAATCTTTATTTCGG	56.1	154	198	061
1.4	71.1	0.04014	TOCCTTTCGTTTATAAACATCC	54.5			
		NGA8 K					
			AUJEAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	50.3	150	140	140
<u>\</u>	102	NGA1107 L	UCGAAAAACAAAAAAAAA				
		NGA1107 R	CGACGAATCGACAGAATTAGG	28			
		0 0000	GAAATCCAAATCCCAGAGGG	58	611	189	611
>	11.8			1.09			
		NGAZZS K	ICICCCACINGINISTICS				
Λ	16.7	NGA249 F	TACCGTCAATTTCATCGCC	55.4	125	115	115
		NGA249 R	GGATCCCTAACTGTAAAATCCC	58.2			
^	19.9	CA72 F	AATCCCAGTAACCAAACACACA	56.3	124	110	110
		CA72 R	CCCAGTCTAACCACGACCAC	61.9			
^	20	NGA151 F	GTTTTGGGAAGTTTTGCTGG	55.8	150	120	130
		NGA151 R	CAGTCTAAAAGCGAGAGTATGATG	58.6			

The state of the s

2000	
77	
ij.	
# .	
įi.	
1	

							-
		0 707 4 501	GTTATEGAGITTICTAGGGCACG	1.09	157	123	1.50
>	24	NGA100 F	OI WICK				
		NGA 106 R	TGCCCCATTTTGTTCTTCTC	55.8			
		Somon	The state of the s				
							133
	0 00	NC 4 130 E	AGAGCTACCAGATCCGATGG	59.9	174	132	132
>_	3/.8	INCALSS I		0 11			
		NGA 139 R	GGTTTCGTTTCACTATCCAGG	55.8			
		W CHION					
							000
		0 201011	GGAGAAAATGTCACTCTCCACC	1.09	231	> 250	300
>_	3	NGA/0 F		0			
		NGA76 R	AGGCATGGGAGACATTTACG	57.8			
		,					
:							
		1 10100112	CTCCACCAATCATGCAAATG	55.8	148	156	146
>	1.19	ATHOOIST L					
		ATHSO191 R	TGATGTTGATGGAGATGGTCA	53.7			
						021	172
7	81.7	NGA129 F	TCAGGAGGAACTAAAGTGAGGG	1.09	//1	117	1
•		NGA129 R	CACACTGAAGATGGTCTTGAGG	60.1			
		i i i i i i i i i i i i i i i i i i i					

and the second of the second s

CLAIMS

- 1. An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.
- 2. A DNA molecule according to claim 1 wherein said polypeptide is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 3. A DNA molecule according to claim 1 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 4. An isolated and purified polypeptide functionally involved in the DNA 10 mismatch repair system of a plant.
 - 5. A polypeptide according to claim 4 which is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 6. An isolated and purified polypeptide selected from the group consisting of a polypeptide encoded by the gene *AtMSH3* (SEQ ID NO: 18), a polypeptide encoded by the gene *AtMSH6* (SEQ ID NO:30), polypeptides homologous to a polypeptide encoded by the gene *AtMSH3* (SEQ ID NO: 18) and polypeptides homologous to a polypeptide encoded by the gene *AtMSH6* (SEQ ID NO:30).
- 7. An isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.
- 8. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence.
- 9. A DNA molecule according to claim 8 wherein said polynucleotide is capable 30 of interfering with the expression of a plant polynucleotide sequence is a sense polynucleotide, an antisense polynucleotide or a ribozyme.
 - 10. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

The second of the second

to the first of the same


according to any one of claims 13-16 and causing said DNA sequence to express said polynucleotide or said polypeptide.

- 23. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a plasmid or vector according to claim 17 and causing said DNA sequence to express said polynucleotide or said polypeptide.
- 24. A process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred.
- 25. A process according to claim 24 wherein a first gene is incapacitated in said first plant, a second gene is incapacitated in said second plant, and said first and second genes are incapacitated in said hybrid plant thereby altering the mismatch repair system of said hybrid plant.
 - 25. A process according to claim 25 wherein said incapacitation of the mismatch repair system of said hybrid plant is reversible.
- 26. A process according to claim 24 wherein a new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait is 20 observable in at least one of said offspring plants.
- 27. A process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.
- 28. A process according to claim 27 wherein said step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene according to claim 13 and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant, cell or cells.
 - 29. A process according to claim 28 comprising inactivating an MSH3 gene and/or an MSH6 gene of said plant.
- 30. A process according to claim 28 comprising inactivating an MSH3 gene and an MSH6 gene of said plant.

- 31. A process according to claim 27 comprising at least partially inactivating the mismatch repair system of said plant in a predetermined cell type or in a predetermined tissue of said plant.
- 32. A process according to claim 31 further comprising restoring mismatch repair 5 in said cell type or said tissue.
 - 33. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule according to claim 1 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
- 34. An oligonucleotide capable of hybridising at 45°C under standard PCR to conditions to the DNA of SEQ ID NO: 18 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
 - 35. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.

The state of the s

Figure 1






Figure 2

Comments/References: 52= 3' side of S5 (AtMSH3) 1244bp in pUC18/Sma1

and the same of th




Figure 3

Comments/References: 13 = 3' side of S5 (AtMSH3) 2104bp in pUC18/Sma1

Figure 4

80	144 15	204 35	264 55	324 75	384 95	444 115	504 135	564 155	624 175	684 195	744 215	804 235	864 255	924
AGGA	CCC	CCG P	CTC L	GAT D	GTT V	AAG K	GGA G	TTC F	GCA A	GCA A	GAA E	AAT N	GAA E	GTT V
ACGG	GCT A	CCA	CAC H	CCC	TAT Y	CTA L	TTC	AAT N	AAT N	GGT G	$_{\rm L}^{\rm CTT}$	AGT S	ATT I	GTT V
TTAA	TTC	CCG	GAC D	GTA V	GAA E	GAG E	${ m TTC}$	CAC H	GTG V	CAT H	ACG T	CAG Q	GGT G	GAA E
TCTC	TTC	ACA T	TCC S	CCA P	GAG E	GTG V	AGA R	GAT D	$_{\rm L}^{\rm CTG}$	TCC	GCC	TCAS	$_{\rm C}^{ m TGT}$	GGT G
CCTAAGAAAGCGCGCGAAAATTGGCAACCCAAGTTCGCCATAGCCACGACCACGACCTTCCATTCTCTTAAACGGAGGA	CGT R	TCA	CTC L	AAC N	CCG P	GTG V	TAC Y	ATG M	AGA R	AAG K	AAA K	GGT G	9 9	ACA
CTTC	TCT	TCA S	$_{\rm L}^{\rm CTT}$	CAA Q	TCG S	CAA Q	AGG R	CAT H	AGA R	ATT I	ACC T	TTT F	TTA L	TCG S
CGAC	ATT I	GAA E	AAG K	ACT T	CCC	CAG Q	TAC Y	GCT A	GTG V	GCC	TAT Y	GG'F G	ACA T	AT'T I
ACCA	ACG	GCC	CGT R	CAC H	gaa E	GAAA E	GGT G	TAC Y	CAT H	GCA A	$_{ m I.}^{ m TTG}$	GAA E	GAG E	GAA E
CACG	CAG Q	GT'A V	AAG K	CCT P	$_{ m L}^{ m CTG}$	${ m TTG}$	GTT V	ATT I	TTC F	ACT T	GCG A	GAA E	TCG S	GTT V
TAGC	CAG Q	CCG P	TCC S	TCT S	TTT F	CCA P	GAA E	GGT G	AAT N	GAA E	TCG S	GGT G	AAG K	ე ე
CCCA	AAG K	AAT N	CCT P	$_{\rm L}^{\rm crr}$	aga R	ACA T	GTG V	rrg	$_{ m L}^{ m TTG}$	ACT T	CTG L	GGT G	GTT V	GTT V
GTTC	CAA Q	ccG P	TCT	AAA K	CAG Q	TAC Y	ATG M	GTG V	CGA R	CAG Q	GCA G	TGT C	AGA R	GTT V
CCAP	AAG K	GAA E	TTC	CCT P	CTC L	AAA K	$_{\rm L}^{\rm TTG}$	CGC R	TTT F	AAG K	CGG R	GGT G	GAG E	GGT G
CAAC	၁၅၅	CAC H	TCC S	AAG K	TTT F	AGG R	GTT V	GCA A	ACA T	GTG V	TTC	GGT G	GAT D	GTC V
TTGG	ATG M	ACT T	GTA V	AAA K	AGA R	TCG S	GTG V	GCA A	CCA P	GTA V	TTT F	AGT S	GTG V	AGA R
AAAA	ATT	CCG P	ACT T	CCC	caa o	TCA S	GAT D	ATC I	GTG V	GGT G	CCT P	ATA I	GTT V	GTT V
ອວອວະ	AGCA	TCC	GCC	TCA	CAC H	TCA	CCA P	GAG E	AGT S	ATT I	9 9	GAT D	TGT C	GAT D
AGCG	GATTACGAATAAAGCAATT	A.A.A K	TCC	GCG A	TTA L	ACG T	TAC Y	GCG A	GCG A	AAG K	ACC	GAG	GTT V	TTT F
AGA	'ACG	CCC	ATA I	GCC		gaa E	AAG K	GAC			CGG R		${ m TTG}$	AGT S
CCTA	GATI	AAA K	AAG K	GCC	CCC	CCC	AGC S	GAA E	ATG M	GGA	AAC	GCG A	TTC	ATG
F	8 7	145 16	205	265 56	325 76	385 96	445 116	505 136	565 156	625 176	665 196	745	805	865 256

the control of the co

2:5:	į
ŧi,	
L.	1
# (E)	į
Łij.	ļ
F	ļ,
	į
W.	
ñ	
	=
Functions:	
Grant's	1
**	ĮĮ.
Pini Com	4
i.	į

98 4 295	1044 315	1104 335	1164 355	1224 375	128 4 395	1344 415	1404 435	1464 455	1524 475	1584 495	1644 515	1704 535	1764 555	1824 575
9.0	H 60	e-1 (*)												
TTG L	GTG V	AGC S	TTA L	GTT V	CAT H	TCA S	AAT N	GTA V	ATA I	TCC	TTT F	GGA G	GCT A	AGG R
AGC 's	TTG	TTC	AAC N	ACA T	TGC	$_{\rm L}^{\rm TTG}$	AAA K	ACA T	TTG L	TCTS	GAG E	CGT R	GAA E	ATG M
TTG	TTT F	TGT C	GGT G	${ m TTG}$	TTT F	TCT	GTG V	CTT L	AAT N	AGT S	CCT P	CAA Q	ATG M	GAA
ATT	AAG K	GAT D	GCA A	TGC	ACG T	CGC R	GTT V	ACA T	AGA R	CAT H	TCA S	ATT I	GTT	S
GTG V	GAG E	CTG L	AGC S	TCT S	CTA L	TTT F	GAG E	CAC H	GAT D	TCT	GTA V	GAT D	GCA	GAC D
GCT A	ACT	TCA	ATC I	ATG M	GCC	TCA S	${ m TTG}$	AAT N	TGC	GGA G	ATT	TCT S	ATT I	AAG CAA K Q nued)
GAG E	CAA Q	GCC	AAA K	GGA G	CTC	GCC	CAG Q	ATG M	CTA L	ATG M	GCA A	TCA	TTC	
TTA L	CAA Q	CGT R	GAA E	AAA K	GCC	999	CAA O	AAT N	CCT P	TGC	AGA R	AGA R	GAG E	ATA I
GGA G	TCA S	gaa E	${f TGT}$	GAA E	CAA	CAA Q	$\frac{\text{CTG}}{\text{L}}$	CAT H	CAT H	GCT A	GAG	TCT S	ACA	966 9 4
AGT S	CTT L	GTG V	TTA L	GCT A	GTT V	TAC Y	ACT T	${ m TTC}$	ACT T	TCT S	TCT	ATG M	, GCC	ccc crr R L
AGA R	CCT P	CGA R	TCA S	GCT A	ACT T	$_{ m L}^{ m CTT}$	AAT N	TTA L	GTG V	ATT I	GGT	GCT	AAA K	: ccc R Fig
ATG M	CAG Q	GTT V	ATT I	GAG E	$_{\rm L}^{\rm CTG}$	ATC	GCC	TCC S	TGG W	GAG E	GAA E	ACA T	GCT	CAG
TTC	299 299	AAC N	GTT V	CTG L	CAT H	AGG R	TCA S	9 299	CAC H	TCT S	GAA E	${ m TTG}$	ACT	ATT I
AAT N	CTT L	TCA S	GAG E	A.A.G K	CCA P	GAA E	CTC	TCT S	AGA R	GTT V	GTT V	GTC	CGG R	CAA Q
GAT D	$_{ m L}^{ m TTG}$	ACC T	GAT D	ATG M	ATG M	TTT F	ACT	GAA E	CTT L	GCT A	${ m r}{ m T}{ m G}$	TCA S	CAT H	AAG K
AAT N	$_{ m L}^{ m CTG}$	CCT P	GTA V	gaa E	AAC N	GGA G	ATG M	TCG	$_{\rm L}^{\rm CTT}$	GAT D	GAG E	TCC	TTT F	9
TTC F	GAG E	GGA G	GCA	AAA K	ATG M	$\mathbf{r}\mathbf{r}\mathbf{r}$	GAG E	GGA G	AGG R	CTT L	AGT S	crc	ATC	GCG A
GAG	GCT	GCT A	AAT N	GAT D	ATT I	CAG Q	ACA T	GAT D	TCC	CGG R			AGA R	CIT
GAA E	CCA P	CAT M	GT	GAT D	ACA T	aaa K	AAC N	TCA S	GCT	GCT A	CTC L	CTC	ACA	TTA L
TAT	TCA		ΑŢ	GAA E	CAT H	CTC L	AGT S	AAT N	TAT	TCT	CAG Q	TAT Y	ATA I	ATT I
925 276			1105 336	1165 356	1225 376	1285 396	1345 416	1405 436	1465 456	1525 476	1585 496	1645 516	1705 536	1765 556

<u>L</u>
M
F
100
F.J
l.i
#
<u> </u>
\$15.2 15.2
₹ ₆
754


																:					
	AGT S	ATG M	CAA Q	TCT S	GCA A	ACT	GTG V	CGA R	TCT	ACT T	CTT	TTG	AGA R	AAA K	TTG . L	ATT	TCT (GTT V	ATT I	TCA S	1884 595
	TCC	CCT P	GTT V	GTG V	GTT V	GAC D	AAT N	GCC	GGA G	AAA K	CTT	CTC	TCT	GCC	CTA	AAT	AAG (GAA	GCG A	GCT A	1944 615
	GTT V	CGA R	GGT G	GAC D	$_{\rm L}^{\rm TTG}$	$_{\rm L}^{\rm CTC}$	GAC D	ATA I	CTA	A'rc I	AC'F T	TCC	AGC s	GAC	CAA	TTT F	CCT	GAG	CTT L	ĠCT A	2004 635
	GAA E	GCT A	CGC R	CAA Q	GCA A	GTT V	TTA L	GTC V	ATC .	AGG R	GAA	AAG	CTG	GAT	TCC S	TCG	ATA I	GCT	TCA S	TTT F	2064 655
	CGC R	AAG K	AAG K	CTC	GCT A	ATT I	CGA R	AAT N	${ m TTG}$	GAA E	TTT	CTT	CAA	GTG V	TCG S	5999	ATC	ACA T	CAT H	$_{ m L}$	2124 675
	ATA I	GAG E	$_{\rm L}^{\rm crg}$	CCC	GTT V	GAT D	TCC	AAG K	GTC V	CCT P	ATG	AAT	TGG W	GTG V	AAA K	GTA	AAT	AGC	ACC	AAG K	2184 695
2	AAG K	ACT T	ATT I	CGA R	TAT Y	CAT H	CCC	CCA P	GAA E	ATA I	GTA V	GCT A	GGG.	T FG L	GAT D	GAG	CTA	GCT A	CTA	GCA A	2244 715
	ACT T	gaa E	CAT H	$_{\mathbf{L}}^{\mathrm{CTT}}$	GCC	ATT I	GTG V	AAC N	CGA R	GCT A	TCG	TGGW	GAT D	AGT S	TTC F	CTC	AAG	AGT	TTC F	AGT S	2304 735
	AGA R	TAC Y	TAC Y	ACA T	GAT D	TTT F	AAG K	GCT A	GCC A	GTT V	CAA	GCT	CTT	GCT A	GCA A	CTG	GAC	TGT C	${ m TTG}$	CAC H	2364 755
	TCC	$c_{ extbf{T}}$	TCA S	ACT T	CTA L	TCT S	AGA R	AAC N	A.A.G K	AAC N	TAT Y	GTC V	CGT R	CCC	GAG E	TTT	GTGV	GAT D	GAC D	rgr C	2424 775
	GAA E	CCA P	GTT V	GAG E	ATA I	AAC N	ATA I	CAG	TCT	GGT G	CGT R	CAT H	CCT P	GTA V	CTG L	GAG	ACT	ATA I	TTA L	CAA Q	2484 795
	GAT D	AAC	${ m TTC}$	GTC V	CCA P	AAT N	GAC D	ACA T	ATT I	${\tt TTG}_{\tt L}$	CAT H	GCA A	GAA E	999	GAA E	TAT	TGC	CAA	ATT I	ATC I	2544 815
	ACC T	GGA G	CCT P	AAC N	ATG M	GGA G	GGA G	AAG K	AGC S	TGC C	TAT Y	ATC	CGT R	CAA Q	GTT V	GCT A	TTA	ATT I	TCC	ATA I	2604 835
	ATG M	GCT A	CAG Q	GTT V	GGT	TCC	TTT F	GTA V	CCA	GCG A	TCA S	TTC	GCC	AAG K	CTG L	CAC H	GTG V	CTT	GAT D	GGT G	2664 855
	GTT V	TTC F	ACT	CGG R	ATG M	GGT G	GCT A	TCA	GAC D	AGT ATC S I	ATC I	cag Q	CAT H onti	NG CAT GGC AG H G R (Continued)	AGA R	AGT	ACC T	TTT F	CTA L	gaa E	2724 875
									•	j P	• }	•		, , ,	ì						

Control of the state of the sta

ĸij.
.IJ
F.
Erroje
N
£.,]

2784 895	2844 915	2904 935	2964 955	3024 975	3084 995	3144	3204 1035	3264 1055	3324 1075	3397 1082	3458 5	3522 16
2725 GAA TTA AGT GAA GCG TCA CAC ATA ATC AGA ACC TGF TCT TCT CGT TCG CTT GTT ATA TTA 2016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1. S E A S H I I R T C S S R S L V I L 8016 F 1.	GAT GAG CIT GGA AGA GGC ACT AGC ACA CAC GAC GGT GTA GCC ATT GCC TAT GCA ACA TTA D E L G R G T S T H D G V A I A Y A T L	CAG CAT CTC CTA GCA GAA AAG AGA TGT TTG GTT CTT TTT GTC ACG CAT TAC CCT GAA ATA O H L L A E K R C L V L F V T H Y P E I	S GCT GAG ATC AGT ATC CCA GGT TCT GTT GGG ACA TAC CAT GTC TCG TAT CTG ACA S GCT GAG ATC AGT AAC GGA TTC CCA GGT TCT GTT GGG ACA TAC CAT GTC TCG TAT CTG ACA	TIG CAG AAG GAT AAA GGC AGT TAT GAT CAT GAT GTG ACC TAC CTA TAT AAG CTT GTG	CGT GGT CTT TGC AGG AGC TTT GGT TTT AAG GTT GCT CAG CTT GCC CAG ATA CCT CCA R G L C S R S F G F K V A Q L A Q I P P	TCA TGT ATA CGT GGC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGA AGA AGA AGA ATG GCT GAG GTA CGT GCA AGA TCA TGT ATA CGT CGT GCA AGA ATG GCT GAG GTA CGT GCA AGA AGA ATG GCT GAG GTA CGT GCT GAG AGA ATG GCT GAG AGA ATG ATG ATG AGA ATG ATG AGA ATG ATG	GAG AGA AAT ACA CGC ATG GGA GAA GGA CAT GAA GAA CCG AGA GGC GCA GAA GAA F R N T R M G E P E G H E E P R G A E E	TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC TCT AIT TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAA TTT GCT CTC TCT GAA GAG GAC GAC	CCT TGG	AAA CCA K P	TACI	GTG TTG CA
0 0	2785 896	2845	, 25	29	3025	3085	3145	3205	32	3,6	က်	34






1 1 1

and the contract

Figure 6






Figure 7

Comments/References: 43= 5' side of S8 (AtMSH6) 2182 bp in pUC18/Sma1

or other comme




Figure 8

Comments/References: 62= 3' side of S8 (AtMSH6) 1379bp in pUC18/Sma1

Ļ	=	j
į,	1	į
1	ij	W.
		ļ
		###
•	5	ļ
	, i	ilens.
	ř	Į
1		
1		Ç.
112	(41)	
The section is a section		
THE LANGE STREET PROPERTY.		then that the
The section of the section of		Kim-Tinn that W
The section of the section of the section of		that there then bear W'
The section of the section of the section of		Kim-Tinn that W

80	153 4	213 24	273 44	333 64	393 84	453 104	513 124	573 144	633 164	693 184	753 204	813 224	873 244	933 264
၁၁၁၃	(n	TCC	999 999	AGA R	SCG P	AAA K	GAT D	AAA K	GAA E	GCT A	GTA V	GAA E	AGA R	TTC
TCC	cAG Q	GTT	GAA (GTT	AAG K	GTA V	AAT N	GGT G	GTA V	CGT R	CCT P	aaa K	AAT N	GTT V
TCTI	s cGC R	TTG (AAG (R	GAG (E	TTT F	TTT	CTG	AAT N	TCA	CCA P	GTT V	AAG K	GCC	GAT D
CCAA	cAG Q	GGT 1 G I	GTG A	GAT (GGA '	AAG K	CCG P	AAT N	AGA R	CGT R	AAG K	GAG E	GAT D	CCT P
ອວອອຸ	ATG M	AAG G K	AAT (GTC (TCT (S	CAT H	GTT (TCC	CTT	ATG M	GAT D	GGA G	AGG R	CCA P
GCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCAAAGGCGAAAATTTTTGGCGCCAATCTTTCCCCCC	CTCTCAGCTCAAAACATCGTTTCTCTCTCACTCTCTCACAATTCCAAAAA	ACG P	TTT F	TCT (000 d	ATG	GTT	CGT R	GAA E	999 9	GAG E	TGT C	ATC I	ATA I
AAAT	TCCA	ACT F	CGA 1	AAA K	CTG (ATT	GAT D	TTT F	GCT A	CCA P	AAG K	GTT V	CGA R	CAC H
GCGA	CAAT	GCG P	CCA (TCG A	GTC (AAT N	GAA	CAA Q	AGA R	ACA T	TTT F	CCG P	TCT	TTA L
CAAG	CTCA	GCG C	GGA (GTT C	CGT (TCC	CGA R	CCT	999	GAA	ACT T	GAT D	TCT	ACC T
GACG	crci	ACC (T	GGA (GCT (CGT R	TTC	AGC S	AT'F I	AGT S	CCA P	ATG M	CAG Q	GAG E	AAG K
CTAC	CACT	CCC F	AGC (S	TTT (F	CCG (CTG	AGG . R	GT'F V	TTC	GGT G	GAA E	CTC L	$_{\rm L}^{\rm crr}$	AGA R
ATTT:	CTCI	AAA C K E	3 299 1 399 1 3	CGT 3	GTT (TCC	GAG	GA'I' D	AGT S	CCT	GAT D	ATG M	TGG W	GAT D
00900	TTCI	CAA F) 299	GTA (AAG K	TCG	GGA G	AAT N	TTT E	GTT V	GAG E	AAA K	gaa E	TAC Y
TTTC	vrcgi	TTC (999	TCT (GAG .	GCT	TCT	GCT A	GCT A	GAT D	$_{\rm L}^{\rm CTG}$	$^{ m CTG}_{ m L}$	TTT F	$_{\rm L}^{\rm ctr}$
ATCG	AAACA	TTC 7) 299	GCT	CCG P	GAT	TGT C	AAG K	CAT H	၁၅၅	GTT V	AGG R	AAA K	CCC
GAGI	TCA	TCT 1	AGC (S	GAC (CCA P	GST G	GAT D	ATG M	AAC N	GAT D	CGA R	A.A.A K	ACC	GAT D
TGAG	CAG	TTG 1	GCT A	3 66C (ACT	GCC	CGA R	TGT C	AGA R	GTA V	AAG K	A.A.C N	GGA G	GAT D
CCC	CTCI	ATT 1		Æ.		TCC	GAT	r,		3A	${ m TTG}$	TCT	GAA E	CCT P
: : : : : : : : : : : : : : : : : : :	AATT	TCG F			ဗ္ဗ	GAA				ATA I	CGC	GAC D	AAC	
а в в в ст т С. В.	TTTCGAATT	AGA T				GCT A					TCT	$_{ m L}^{ m r}$	GTA	AGA R
-										634	694 185	754	814	874 245

ij.
L
IJ
IJ
ű.
ž:
-1

993 284	1053 304	1113 324	1173 344	1233 364	1293 384	1353 404	1413 424	1473 444	1533 464	1593 484	1653 504	1713 524	1773 544	1833 564
ATT I	4 GGT G	r ggt g	A GTT V	T ATA I	ງ ອອງ ວ	T TCA S	C ATC I	G GAA E	T ACG T	A GAT D	A TCA S	A GAG E	T TTT F	T GAG E
GAC D	TTA L	GTT V	AAA K	ACT T	ATC I	TGT C	TCC	AAG K	TAT	ACA	GAA E	GGA G	ATT	. CTT L
ATG (M	gaa E	CAG Q	TAT Y	AAT N	AAC N	AAG K	9 9	CCA P	aaa K	GAT D	TCT S	CTT L	GAT D	AAT N
TAT A Y	GCG A	AGA R	GGA G	GCT	GGA G	CAA Q	GTT V	TCT S	AGG R	999	TCT	GCT A	999 9	GTA V
GAA TE	GAT D	TGC	CGT R	GGT G	GAG E	CTA L	TGG W	$_{\rm V}^{\rm GTT}$	CTA L	ATG M	GGT G	AGT S	CAT H	ATG M
AGT G S	CTA L	AAA K	GCT A	AGA R	AGC S	GAG E	TTT F	CAG Q	GCT A	GTA V	A.N.A K	$_{\rm L}^{\rm CTT}$	AAG K	ACG T
AAG A K	GAG E	GGA G	$_{ m L}^{ m TTA}$	GCC	GCA A	ATG M	AGG R	ATG M	AAG K	CAA	TTT F	GCC A	$_{\rm L}^{\rm CTT}$	CAG Q .nued
GTT A V	TAT Y	GTG V	CTA L	AAA K	ACA T	AAA K	TTG L	${ m TTG}$	CAA Q	CCA P	TAC Y	GTT V	GTA V	GGC G
AGT G S	$^{ m CTG}_{ m L}$	GGT G	AAG K	GCA A	TCA S	ATC	GCC	TTA L	GCA A	GTA V	GGA G	GAT D	GAT D	GAT D 9 (C
TGG A W	GAG E	AGT S	CAA Q	CAA Q	CCA P	GAG E	GCT	GCG A	GAA E	CCA P	AAC N	TGT	GAA E	A ATT I gure
TAT T Y	TAT Y	ATG M	GTG V	GAC D	ACT T	AAA K	TGT	GGA G	AGA R	GCT	TCT S	GAA E	CTA L	AGA R Fig
CAA T Q	TTT	ACC	GCA A	TCT S	TTA L	ATA I	GAC D	CTT L	TCA S	TTG L	GAA E	AAT N	AAG K	CTC L
AAG C K	aaa K	ATG M	GAG E	ACA T	GTA V	GCT A	GTT V	GCT A	CTA L	CAG Q	ATA I	CTA L	CTA L	rgr C
CAA A Q	999	AAG K	GAT D	GAA E	CAG Q	$_{\rm L}^{\rm crT}$	TTT F	GCT A	999 9	GTA V	ATA I	GGT G	AGG R	GGT G
rca c	GTG V	7GG W	ATA I	CTA L	GTT V	$_{\rm L}^{\rm CTT}$	GCT A	$_{ m C}^{ m TGT}$	AAA K	GCG A	AAT N	GAT D	TCT S	AGG R
GCA T	AAA K	GAC D		CAG Q	CTA L	CAT H	TTT F	TCA S	AGT S	ACG T	AGA R	GTT V	$_{\rm L}^{\rm cTG}$	TAC Y
TCT G	TTT F	$_{\rm L}^{\rm CTT}$	AGT S	GAG E	AAG K	GTC V	GGA G	GCA A	GAC D	TCT S	GTT V	GCT A	CAT H	GTT V
	TTC F	GAG E	GAA E	ATC I	AGG R	GCC	TAT Y	GAT D	TAT Y		GGA G	TGT C	AAT N	CAA Q
AAA ATG K M	$_{\rm L}^{\rm cTT}$	AAG K	TCT S	CGA R	CCA P	GAT D	GTG V	GAT D	TTA L	ACA T	GCT A	AAC	ATT I	TAC Y
AAG A K	GTG V	CAC H	ATC	GGA G	ATT I	CCT	ACT	AGC S	GTG V	${ m TTG}$	GCT A	TGG W	CTA L	CCA P
934 265	996 285	1054 305	1114 325	1174 345	1234 365	1294 385	1354 405	1414 425	1474 445	1534 465	1594 485	1654 505	1714 525	1774 545

The state of the fill of the state of the st

7
LF1
T.
ų.
Ħ
Li
121
E
ļai.
٠
F===
Carles to be


1893 584	1953 604	2013 624	2073 644	2133 664	2193 684	2253 704	2313 724	2373 744	2433 764	2493 784	2553 804	2613 824	2673 844	2733
AAC N	GAT D	AGT S	CGC R	. GTG V	GAT	, AAA K	22C	ACT T	CAC	GGA	A ACA T	r ggr g	S ATT	r cgr R
GAT D	AAA K	gaa E	GGA G	AAA K	ATT I	TGT C	GCA A	gaa E	ATT I	GCT A	AAA K	GAT D	AGC	CTT
CTT L	CTC L	TCA S	CTC L	AAA K	GGA G	CTC L	gaa E	GCT A	GTC V	TCT	CAG Q	GCC	9 299	CTT
TAT Y	CCA P	AAC N	$\mathop{\mathrm{CTG}}_{\mathrm{L}}$	999 9	AGT S	AAA K	$_{ m F}$	AAC N	GAG E	CTC L	aat n	GCA A	AGT S	ACT
AAA K	CAT H	GCA A	AGA R	CTG L	AGA R	TAT Y	CAA Q	gaa E	TCT	AGT S	CAG Q	GTT V	AGC S	TCAS
TAC Y	TGC	ACG T	GAA E	C'IT L	TTC F	CTT L	TCT S	GAT D	TGG W	GCA A	GAT D	GCA	AGA R	AAA K
TTG L	ATC I	TTC F	TTA L	GCT	9	TTG L	CTT L	ACA T	CAA	GCA A	ACA T	TTT F	AGA R	GGA 1 G 1
ACC	7GG ¥	GAA E	GAC D	CCT P	AAA K	AGT S	TTT F	GTG V	ACT T	ATC I	GCT A	CCA P	GCT A	G GGC G (Conti
999	AAT N	GAA E	CCA P	TTG L	GTG V	ATG M	TTA L	GAT D	GCA A	GCA A	GAA E	CAT H	GAG E	M 6
TCA	AGG R	GTT V	CTT L	GTG V	ATT 1	ATG M	GAG E	CAA Q	AGA R	TTT F	TCA S	TGG W	199	AAC N pure
CCT	T'TA L	GTA V	AAA K	TCT S	CAA	AAT N	CTA L	AAC	GAA	TCT	GAA E	CTA L	$_{\rm L}^{\rm CTT}$	CCA P Fig
GGT	CTC L	GAT D	CAC H	GCC	999	TCA	9 999	CAG Q	ATC 1	AGA R	CCC	GGA G	CTC	GGA G
GGT	CGA R	CTT	CTC L	TCA	ፐፐፐ F	GAA E	AGC S	TAT Y	TTT F	CTG L	TTT F	CAA Q	ATA I	ACG T
GAT (A.A.G. K	CGG R	TAT Y	TCA S	GCA A	A.A.G K	aaa K	AAT N	CTT L	GTC V	ATT I	ATC I	GAT D	$^{ m CTG}_{ m L}$
rgr (GGT G	AAA K	CAG	CGA R	AAA K	CAG Q	GGA G	CCA P	gaa E	GAT D	GTT V	AAA K	AAT N	TTA L
AGC .	ACT	AAT	299	GTT V	GTT V	CTA L	GTA V	TTT F	ATC I	CTA L	CCT	$_{\rm L}^{\rm crr}$	CCG P	${ m TTG}$
AAT N	CCA J	ATC	ACT	AGC S	CGA R	GCT A	TTA L	GAC D	$_{\rm L}^{\rm crr}$	TGC C	AGG R	ATA I	GTT V	TCAS
AAC 1	AGT (AGC S	ATC	TCT	CAA Q	$_{ m L}$	ATA I	AGC S	ATA I	AGC S	GCC	CCA	CCT	CGG R
TTT A	GTT 1	GAA 1	CAA	AAG K	AAA K	TTG	CCT	GAT D	ACA T	ATA I	ATG M	999 999	${ m TTG}$	CCT P
ATA 1	TGT C	GTA (ATG (ATC I	CTG L	CTG	CTT	ATA I	CTC	ACC	AGC	AAA K	CAA	CAT H
1834 565	1894 585	1954 605	2014 625	2074	2134 665	2194 685	2254 705	2314	2374	2434 765	2494 785	2554 805	2614 825	2674 845

Ť,	7
1	M
1	Į
1	
il de	Ų
1	ij
ı,	N
15	
_	
1	
mile that then	
mile that then	7
mile that then all	

2793 884	2853 904	2913 924	2973 944	3033 964	3093 984	3153 1004	3213 1024	3273 1044	3333 1064	3393	3453 1104	3521 5	3579	3606 28
7GC C	ACA T	ACT T	TAC Y	TTT F			GTG V	A A	A AGT S	r GCC	S TCT	CACTATCTGAAGCTCGTTAAGTCTTTTGCCTCTCT	aaa K	
TCT S	ATG M	GCA A	GGA G	CTC L	TCG S	CAA O	CAA Q	CAA Q	TCA S	AAT N	TCC S	CCC	TTA L	
GAG	ATC	AAT N	GAT D	ATG M	ACC T	GAT D	CTT L	GCT A	TTC	AAC N	A.A.A K	TTT	CGA 1 R 1	
TGT (AGA R	CAG	TTC F	CGG R	GTC V	TGT C	GGA G	GCT A	GAG E	CAC H	ATC I	AGTC	CCT C	
CCG 7	GAT D	CTT	ACT T	TGT C	CGT R	GGT G	TAC Y	GGT G	TCT S	GCC A	GAG E	GTTA	TTT C F P	
GTG (TCT	GTT V	AGT S	CAA Q	CCA P	CGT R	AGC S	TCA S	AGA R	GTC V	CAT H	GCTC	TTG T	ਰੇ
TAC Y	GCA A	TCA S	ACT T	GTT V	CAC	CCA P	GAG E	GCA A	C'I'A L	CGA R	TGG W	TGAA	AAA T K	(Continued)
TGC	၁၁၅	GCG A	GGA G	AAA K	TCT	CAA Q	CCT P	ACA T	GAG E	TCT	TTA L	TATC	AAA P K	Cont
၁၅၅	CTT L	ACA T	AGA R	GAG E	GCG	TAT Y	${ m TGT}$	GAA E	AGT S	ATT I	TGC	CAC	ATC P	6
CTT L	AGG R	GAG E	ე ეეე	GTA V	TTC	GAT D	GCT A	GT'ľ V	TCA S	GGT	TTT F	TAA *	TAT A	Figure
CAA Q	ACA T	ACT T	CTG L	CTG L	GAA E	TCT	GGA G	GTG V	AAG K	GTG V	$\mathop{\mathrm{TTG}}_{\mathrm{L}}$	ACA T	TTA T	
GCC	${ m TTC}$	TGC	GAA E	CAC H	AAG K	AGA R	GAG E	CAA Q	TTC F	${ t TTG}$	ACT	ATG M	TGC T	A AAA K
TTT F	ATC I	GAA E	GAC D	CGT R	ACC T	TCA S	ACC T	AAC N	AAC	TCA S	GAC	GCT A	AAA T	A AAA K
ATC	ACT T	GTA V	$_{\rm L}^{\rm crr}$	FTT	CTC L	AAA K	\mathtt{TTA}	CCA P	GGA E	AAG K	TAC	, ATG M	AAA P K	A AAA K
GTT	GAT D	${ m TTG}_{ m L}$	ATC I	GTT V	CCT P	TTC F	CGT R	ATA I	999 9	$_{ m L}^{ m CTC}$	GAC D	TAA *	CTT P	A AAA K
GCC A	GTG V	TTT F	GTA V	TCG	CAC H	GCA A	TAC Y	GGA G	A'l''F I		GAT	AAA K	CCT (A AAA K
CTG	CTC	ACC	CTA	TAC	TAC Y	TGC C	$\mathop{\mathrm{TTG}}_{\mathrm{L}}$	GCT A			GAA E	CCC	ATT C	A AAA K
TGT	()		TCA	GCA	CAT H	GCT A	${ m TTC}$						TTT P	A AAA K
ACA			GAT	ATT I	ACA T	ATG M	GTG V						ATG T	AAA K
GCA	GAA					CAC H	CTA	GCA	ATG	CTG L	CCC	TAC	α, Σ υ	aaa K
2734	2794 885	2854 905	2914 2914	2974 945	3034 965	3094 985	3154	3214 1025	3274	3334 1065	3394	3454	3522	3580 20

the state of the s

Figure 10



TTTTTTGGTTGCTAACAATAAAGGTATACGGTTTTATGTCATCAATATAA	50
CTATATATAAAAGAAATGAAAGATATATATTGTTTTTTCATTTATCAAAC	100
AAAACAACAAGACTTTTTTTTTTTTTTTTTTTTTTTTTT	150
GATAAACGACATCGTTTAATCATTTCCCAATTTTACCCCTAAGTTTAACA	200
CCTAGAACCTTCTCCATCTTCGCAAGCACAGCCTGATTAGGAACAGCTTT	250
ACCATTCTCATATTCCTGAACTACCTGAGTCCTCTCATTGATCTGTTTCG	300
CCAAATCCGCTTGTGACATCTTCTTCTCCAATCTCGCTTTCTGTATCATC	350
AACCTCACCTCTGCTTTCACACGATCCATCGCCGCAGGCTCTGTTTCTTC	400
TTCCAGCTTCTTCGTGTTAATCACCGGAACCGCCGTAGATTTCCCCTTTT	450
TGTTCGAACCGGCATCGAATTTCTTAACCGTTTGAACCGCGACACCGTTT	500
CTCAGAGCTGCGTTAACCGCTTTCGGATCGCGTAGGTCTTGGCTCTTTTG	550
TTTTGATTTGTGGAGAACTACTGGTTCCCAGTCTTGTGTTACTGCTCCTG	600
GGTATCTGCTCGGCATCGTCGATGAATTGAGAGAAAGGAACAACGCGAAA	650
ATTTTATTAATCTGAGTTTTGAAATTGAGAAACGATGAAGATGAAGAATG	700
TTGTTGAGAGGATTGTGATATTTATATATACGAAGATTGGTTTCTGGAGA	750
ATTCGATCATCTTTTCTCCATTTTCGTCTCTGGAACGTTCTTAGAGATG	800
ATTGACGACGTGTCATTATCTGATTTGCAGTTAACCAATGCTTTTTGGGT	850
TGGATTCGTGGTACACCATATTATCCGATTTGGCTCAATGGTTTTATATA	900
AATTTGGTTTTCGGTTCGGTTATGAGTTATCATTAAAATTAAGCTAACCA	950
AAAATTTTCGTAAAATTTATTTCGGTTTCAATTCGGATCCCTTACTTCCA	1000
GAACCGAATTATTCGAAACCGGGGTTAGCCGAACCGAATACCAATGCCTG	1050
ATTGACTCGTTGGCTAGAAAGATCCAACGGTATACAATAATAGAACATAA	1100
ATCGGACGGTCATCAAAGCCTCAAAGAGTGAACAGTCAACAAAAAAGTT	1150
GAGCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCAAGGCGAAAAT	1200
TTTTGGCGCCAATCTTTCCCCCCCTTTCGAATTCTCTCAGCTCAAAACATC	1250
GTTTCTCTCTCACTCTCTCACAATTCCAAAAAATGCAGCGCCAGAGAT	1300
CGATTTTGTCTTTCCAAAAACCCACGGCGGCGACTACGAAGGGTTTG	1350
GTTTCCGGCGATGCTGCTAGCGGCGGGGGGGGGCGGCGGAGACCACGATTT	1400
AATGTGAAGGAAGGGGATGCTAAAGGCGACGCTTCTGTACGTTTTGCTGT	1450
TTCGAAATCTGTCGATGAGGTTAGAGGAACGGATACTCCACCGGAGAAGG	1500
TTCCGCGTCGTGTCCTGCGTCTGGATTTAAGCCGGCTGAATCCGCCGGT	1550
GATGCTTCGTCCCTGTTCTCCAATATTATGCATAAGTTTGTAAAAGTCGA	1600
TGATCGAGATTGTTCTGGAGAGAGGTACTAATCTTCGATTCTCTTAATTT	1650
TGTTATCTTTAGCTGGAAGAAGAAGATTCGTGTAATTTGTTGTATTCGTT	1700
GGAGAGATTCTGATTACTGCATTGGATCGTTGTTTACAAATTTTCAGGAG	1750
CCGAGAAGATGTTGTTCCGCTGAATGATTCATCTCTATGTATG	1800
ATGATGTTATTCCTCAATTTCGTTCCAATAATGGTAAAACTCAAGAAAGA	1850
AACCATGCTTTTAGTTTCAGTGGGAGAGCTGAACTTAGATCAGTAGAAGA	1900
TATAGGAGTAGATGGCGATGTTCCTGGTCCAGAAACACCAGGGATGCGTC	1950
CACGTGCTTCTCGCTTGAAGCGAGTTCTGGAGGATGAAATGACTTTTAAG	2000
GAGGATAAGGTTCCTGTATTGGACTCTAACAAAAGGCTGAAAATGCTCCA	2050
GGATCCGGTTTGTGGAGAGAAGAAGAAGTAAACGAAGGAACCAAATTTG	2100
AATGGCTTGAGTCTTCTCGAATCAGGGGATGCCAATAGAAGACGTCCTGAT	2150
GATCCCCTTTACGATAGAAAGACCTTACACATACCACCTGATGTTTTCAA	2200

GAAAATGTCTGCATCACAAAAGCAATATTGGAGTGTTAAGAGTGAATATA	2250
TGGACATTGTGCTTTTCTTTAAAGTGGTTAGTAACTATTAATCTAGTGTT	2300
CAATCCATTTCCTCAATGTGATTTGTTCACTTACATCTGTTTACGTTATG	2350
CTCTTCTCAGGGGAAATTTTATGAGCTGTATGAGCTAGATGCGGAATTAG	2400
GTCACAAGGAGCTTGACTGGAAGATGACCATGAGTGGTGTGGGAAAATGC	2450
AGACAGGTAAATTAGTTGAAACAACTGGCCTGCTTGAATTATTGTGTCTA	2500
TAAATTTTGACACCACCTTTTGTTTCAGGTTGGTATCTCTGAAAGTGGGA	2550
TAGATGAGGCAGTGCAAAAGCTATTAGCTCGTGGGTAAGGGAACCATCAT	2600
ACTTTATGGAATTCGTTTACTGCTACTTCGGCTAGGATTTAAGAAATGGA	2650
AATCACTTCAAGCATCATTAGTTAGGATCCTGAGAACTCAGGATGTTTTC	2700
TTATTCGTTATATAATAAGTCTTTTCATCAAGGAGTAACAAACA	2750
GCACAATATTTGTGTGCTCACTGGCAAGGCATATATACCCAGCTAACCTT	2800
TGCTAGTTCACTGTAGTAACAGTTACGGATAATATATGTTTACTTGTATG	2850
TGGTACCCTCATTTGTCTCTCATGGAGGCTTTGAAGCCTTGTGTTGAAA	2900
CTGGATAGTTACATATGCTTCCAACAGAAACTAGCATGCAGATTCATATG	2950
CTTTCCTATTCTACTAATTATGTATTGACACACTCGTTGTTTCTTTTGAA	3000
AGATATAAAGTTGGACGAATCGAGCAGCTAGAAACATCTGACCAAGCAAA	3050
AGCCAGAGGTGCTAATACTGTAAGTTTTCTTGGATAGGTCAAGGAGAGTG	3100
TTGCAGACTGTTTTTGATCATTTCTTTTTTTTTGTACATTACTTTCATGCTG	3150
TAATTAACTCAATGGCTATTCTGGTCTGATTATCAGATAATTCCAAGGAA	3200
GCTAGTTCAGGTATTAACTCCATCAACAGCAAGCGAGGGAAACATCGGGC	3250
CTGATGCCGTCCATCTTCTTGCTATAAAAGAGGTTTGTTATTTACTTATT	3300
TATCTTATCATGTTCAGTTCATCCAAGTCCTGAAAAATTACACTCTTCTT	3350
TACCAATCTTCCATCAAGCTGTGTAAAGGATTTGGAATTAGAAAATCATT	3400
ATTTGATGCTTTGTTTTATATGCAAGAGGTTCCCTTGAAAAGATCTGTTT	3450
AAGATTCTTTGCACTTGAAAAATTCAATCTTTTTAAGTGAATCCCCTACT	3500
TTCTTACAATGATCATAGTCTGCAATTGCATGTCAAGTAATATCATTCCT	3550
TGTTACTGCATCCCCCTCTTTCTTAATGACCATTGTCTATGTTGTTTTT	3600
TCTCGTGTGCTGGAGAAAATGATAGCTGATCCAAGCTGTACATTATCATG	3650
ATTAAGTAGCTGCTCAGGAATTGCCTTTGGTTACATTGCCTAATGGTTTG	3700
ATGTCAATTTTTCTTCTGAATCTTTATTTTAGATCAAAATGGAGCTACAA	3750
AAGTGTTCAACTGTGTATGGATTTGCTTTTGTTGACTGTGCTGCCTTGAG	3800
GTTTTGGGTTGGGTCCATCAGCGATGATGCATCATGTGCTGCTCTTGGAG	3850
CGTTATTGATGCAGGTAAGCAAGTGTATTCTGTATCTTATGTGTACCATG	3900
TGACTTCCTGTGCATATATTTGGGTTGCAGGAACTAATTCTGAATCACCA	3950
TTTGGTATGTTTTTCCAGGTTTCTCCAAAGGAAGTGTTATATGACAGTA	4000
AAGGTAAACTGCTTGTATCGCCAGTTGTTTTGTTAAACAGAATTTAAGGT	4050
AAATGACACTGGTTAATTTAAAGTGCATACATGTTGAAATATTGCAGGGC	4100
TATCAAGAGAAGCACAAAAGGCTCTAAGGAAATATACGTTGACAGGTACC	4150
ATTTCAGTAGGCAAGCTAACTGACAATTTAACCGCTCACCGAATGATAGG	4200
TCTCTTAAACATTGCTAATGTAGATGATGTTTATGTTTCAATCTAATAGG	4250
GTCTACGGCGGTACAGTTGGCTCCAGTACCACAAGTAATGGGGGGATACAG	4300
ATGCTGCTGGAGTTAGAAATATAATAGAATCTAACGGATACTTTAAAGGT	4350
TCTTCTGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATG	4400
TOTICIGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATGTGATGT	1100

	*
TGCCCTTAGTGCTCTTGGAGAGCTAATTAATCATCTGTCTAGGCTAAAGG	4450
TGTGTTGGCTTGTTTAGTTTTTGCTTTTCACAAATTAAGCAAAGGAACTT	4500
TTCATAACTTACAGTTTCTATCTACTTGCAGCTAGAAGATGTACTTAAGC	4550
ATGGGGATATTTTCCATACCAAGTTTACAGGGGTTGTCTCAGAATTGAT	4600
GGCCAGACGATGGTAAATCTTGAGATATTTAACAATAGCTGTGATGGTGG	4650
TCCTTCAGGCAAGTGCATATTTCTTTTTTGATAACTTCAACTAGAGGGCA	4700
GACATAGAAGGAAAAATTCTAATACTTCGTACGGATCTCCAGTAAGTA	4750
AGCCGATTTTTGTTTACCTATGTAGGGACCTTGTACAAATATCTTGATAA	4800
CTGTGTTAGTCCAACTGGTAAGCGACTCTTAAGGAATTGGATCTGCCATC	4850
CACTCAAAGATGTAGAAAGCATCAATAAACGGCTTGATGTAGTTGAAGAA	4900
TTCACGGCAAACTCAGAAAGTATGCAAATCACTGGCCAGTATCTCCACAA	4950
ACTTCCAGACTTAGAAAGACTGCTCGGACGCATCAAGTCTAGCGTTCGAT	5000
CATCAGCCTCTGTTGCCTGCTCTTCTGGGGAAAAAAGTGCTGAAACAA	5050
CGAGTAAGTATCAATCACAAGTTTTCTGAGTAATGCCTTCCATGAGTAGT	5100
ATAGGACTAAAACATTACGGGTCTAGCTAAAGACTGTTCTCCTTCTTTTG	5150
CAATGTCTGGTTATTCATTACATTTCTCTTAACTTATTGCATTGCAGGTT	5200
AAAGCATTTGGGCAAATTGTGAAAGGGTTCAGAAGTGGAATTGATCTGTT	5250
GTTGGCTCTACAGAAGGAATCAAATATGATGAGTTTGCTTTATAAACTCT	5300
GTAAACTTCCTATATTAGTAGGAAAAAGCGGGCTAGAGTTATTTCTTTC	5350
CAATTCGAAGCAGCCATAGATAGCGACTTTCCAAATTATCAGGTGCCCAT	5400
CTATCTTTCATACTTTACAACAAAATGTCTGTCACTACTCAAAGCAATGC	5450
ATATGGCTTAGATCTCAACTCACACCCCGAGGATCCTAAAGGGATTTGCT	5500
TTTTATTCCTAATGTTTTTGGATGGTTTGATTTATTTCTAACTTGAACTT	5550
ATTAATCTTGTACCAGAACCAAGATGTGACAGATGAAAACGCTGAAACTC	5600
TCACAATACTTATCGAACTTTTTATCGAAAGAGCAACTCAATGGTCTGAG	5 65 0
GTCATTCACACCATAAGCTGCCTAGATGTCCTGAGATCTTTTGCAATCGC	5700
AGCAAGTCTCTCTGCTGGAAGCATGGCCAGGCCTGTTATTTTTCCCGAAT	5750
CAGAAGCTACAGATCAGAATCAGAAAACAAAAGGGCCAATACTTAAAATC	5800
CAAGGACTATGGCATCCATTTGCAGTTGCAGCCGATGGTCAATTGCCTGT	5850
TCCGAATGATATACTCCTTGGCGAGGCTAGAAGAAGCAGTGGCAGCATTC	5900
ATCCTCGGTCATTGTTACTGACGGGACCAAACATGGGCGGAAAATCAACT	5950
CTTCTTCGTGCAACATGTCTGGCCGTTATCTTTGCCCAAGTTTGTATACT	6000
CGTTAGATAATTACTCTATTCTTTGCAATCAGTTCTTCAACATGAATAAT	6050
AAATTCTGTTTTCTGTCTGCAGCTTGGCTGCTACGTGCCGTGTGAGTCTT	6100
GCGAAATCTCCCTCGTGGATACTATCTTCACAAGGCTTGGCGCATCTGAT	6150
AGAATCATGACAGGAGAGAGTAAGTTTTGTTCTCAAAATACCAATTCCTC	6200
GAACTATTTACTCAGATTTGTCTGATTGGACAAGGTGGTTTTGCTTTTT	6250
TTTAGGTACCTTTTTGGTAGAATGCACTGAGACAGCGTCAGTTCTTCAGA	6300
ATGCAACTCAGGATTCACTAGTAATCCTTGACGAACTGGGCAGAGGAACT	6350
AGTACTTTCGATGGATACGCCATTGCATACTCGGTAACCTGCTCTTCTCC	6400
TTCAACTTATACTTGTTGATCAACAAAAACATGCAATTCATTTTGCTGAA	6450
ACTTATTGATTTATATCAGGTTTTTCGTCACCTGGTAGAGAAAGTTCAAT	6500
GTCGGATGCTCTTTGCAACACATTACCACCCTCTCACCAAGGAATTCGCG	6550
TCTCACCCACGTGTCACCTCGAAACACATGGCTTGCGCATTCAAATCAAG	6600

Figure 11 (Continued)

THE PROOF STATE

 $(0,1) = (0,1) \cdot (0,1$

6650 ATCTGATTATCAACCACGTGGTTGTGATCAAGACCTAGTGTTCTTGTACC 6700 GTTTAACCGAGGGAGCTTGTCCTGAGAGCTACGGACTTCAAGTGGCACTC 6750 ATGGCTGGAATACCAAACCAAGTGGTTGAAACAGCATCAGGTGCTGCTCA 6800 6850 CTGAGTTCTCAAGTCTGCATGAAGACTGGCTCAAGTCATTGGTGGGTATT 6900 TCTCGAGTCGCCCACAACAATGCCCCCATTGGCGAAGATGACTACGACAC 6950 TTTGTTTTGCTTATGGCATGAGATCAAATCCTCTTACTGTGTTCCCAAAT 7000 AAATGGCTATGACATAACACTATCTGAAGCTCGTTAAGTCTTTTGCTTCT 7050 CTGATGTTTATTCCTCTTAAAAAATGCTTATATATCAAAAAATTGTTTCC 7100 TCGATTATAACAAGATTATATATGTATCTGTCGGTTTAGCTATGGTATAT 7150 AATATATGTATGTTCATGAGATTGGTCAAGAGAAATACTCACAAACAGTA 7200 TATTAAGAAGGAAATATGTTTATGCATTAATTTAAGTTTCAAGATAAACT 7250 GCAAATAACCTCGACTAAAGTTGCAAAGACCAAACACAAATTACAAAACT 7300 7350 TATTTTGTTGCATCTACAAACAACACAAACCTACATAGTTTATAACTTAC 7400 TCATCACTGAGATTAACATCAGAATCATTCTCCATTTCTTCATCTTCACT 7450 CTCATCATCACCACCACCATGATGATTCTCCTCCTCTTCACGTAACC 7500 TAGCAATCTCACTCTGAGCTCTATCAACAATCTGCTTCTTCTGCAACTCC 7550 AAATCTCTCTGAAAATCAGCTCTCATCTTCTCCAACTCCTTCATTTGCTC 7600 TTTCTTACTCTTCTCCATCTTCTCATAAACCTTCCCAAACCTCTCAACAG 7650 AATCCGCCAACATCTTATACGAAGCAGCGTCATTAACCTTCTTCCTCTCG 7700 TACTCAACCTCATCCTCATCCTCCTCCTCTCAGAATCACCAGGACT 7750 7800 TCATAAACACAAACTCACCTGAATCAACACCACAAGCTAAACCTAAATCC 7850 GACTTGGGCGAAACACAAAGCAACATATCCAACTTATTGAAAAACGACCA 7900 TTTACTTGAACCTAAACCTGATTTCTCAACCTTAATCTTCTCTTTTCTAT 7950 ACTTCCTCTTCAAGTCATCAATCATTCTCCTACATTGCGTCTCAGATTTC 8000 TCCATCCTTAGCTCCTCACTCACTTCTCAGCTACTTCATTCCAATCCTC 8050 GTTCCTCAAACTCCTTCTACCCAATTGCAAAAACCTATCTCCCCAAACTT 8062 CAAGCAACACAA

Figure 11 (Continued)

The state of the state of

1 1




Figure 12

Comments/References: AtMSH3 3' side antisense : AtMSH3 3' (13 = 2104bp) from pUC18/13 Sal1/Sst1/T4 into pCW164 BamH1/T4 in Agrobacterium LBA44O4




Figure 13

Comments/References: AtMSH6 (S8) 3' side antisens : 62 Sal1/Sst1/T4 (1379bp) into pCW164 BamH1/T4




Figure 14

Comments/References: AtMSH6 3'/AtMSH3 3' antisense : AtMSH6 (S8) 3' side (62=1379bp) Sal1/Sst1/T4 into pPF13 (pCW164 AtMSH3 (S5) 3' side (13=2104) antisens)/Sma1. in LBA4404




Figure 15

Comments/References: AtMSH3 3'/AtMSH6 3' antisens (D): AtMSH3 (S5) 3' side (13=2104bp) Sal1/Sst1/T4 into pPF14 (AtMSH6 (S8) 3'side (62=1379bp) antisense into pCW164)/Sma1. in LBA4404




Figure 16

Comments/References: AtMSH3 (S8) complete, sense orientation: pPF26 (3342bp)

Sma1 into pCW164 Sma1





Figure 17

Comments/References: pPZP111 with codA EcoR1 cassette in EcoR1 site and HInd3 GUS cassette in Hind3 site. KanR. All genes under Promoter/terminator 35S

Figure 18




THE REPORT OF THE PARTY.

1.00

Figure 19

p3243



1.11

1

SEQUENCE LISTING

	<110>	Rhone-Poulenc Agro; Betzner, Andreas Stefan; Doutriaux, Marie-Pascale; Freyssinet, Georges; Perez, Pascual.
ب	<120>	Methods for obtaining plant varieties
š	<130>	395498C
	<150>	PO9745
	<151>	1997-10-10
	<160>	98
post.	<210>	1
	<211>	23
1545 1545	<212>	DNA
The first face will find the first face will be for the face will be for the first face will be for the first face will be for the face will be face will be for the face will be for the face will be for the face will be face will b	<213>	Artificial sequence
	<220>	
n.	<221>	modified_base
	<222>	11
111	<223>	I
H	<223>	<u> </u>
Emilia Emilia	<220>	
	<221>	modified_base
T.	<222>	14
F	<223>	I
	\2.00°	
Taxanii I	<220>	
	<221>	modified_base
	<222>	17
		I
	<223>	±
	<220>	and
	<223>	Degenerate oligonucleotides UPMU used to isolate AtMSH3 and AtMSH6.
	<300>	
	<301>	Reenan and Kolodner
	<302>	Genetics
ž	<303>	132
	<306>	963-973
	<307>	1992
-	<400>	1
	ctggatcca	ac nggnccnaay atg 23

and the second s

<210> 2 <211> 23 <212> DNA WO 99/19492 PCT/EP98/06977

2

```
Artificial sequence
    <213>
    <220>
    <221>
                  modified_base
    <222>
                  15
    <223>
                  I
    <220>
                  modified base
    <221>
    <222>
                  18
    <223>
                  I
    <220>
    <223>
                  Degenerate oligonucleotides DOMU used to isolate AtMSH3 and
                  AtMSH6.
    <300>
hand them then the transfer they form
    <301>
                  Reenan and Kolodner
    <302>
                  Genetics
    <303>
                  132
    <306>
                  963-973
    <307>
                  1992
    <400>
                  2
Ē
with the time that the property
    ctggatccrt artgngtnrc raa
                                                                                 23
                   3
     <210>
     <211>
                  24
     <212>
                  DNA
     <213>
                  Artificial sequence
     <220>
     <223>
                   MSH3 specific primer 636 for PCR using cDNA of Arabidopsis
                   thaliana ecotype Columbia
     <400>
                   3
    tgctagtgcc tcttgcaagc tcat
                                                                                 24
    <210>
     <211>
                   27
     <212>
                   DNA
     <213>
                   Artificial sequence
     <220>
     <223>
                   Primer AP1 for PCR using cDNA of Arabidopsis thaliana ecotype
                   Columbia containing adapter sequences ligated to both its
                   ends
     <400>
                   4
```

. The state of th

Fris
113
H
Fil.
Ħ
! =4
Terror
fij
1

	ccatcctaat a	acgactcact atagggc	27
	<210>	5	
	<211>	23	
5	<212>	DNA	
	<213>	Artificial sequence	
•	<220>		
	<223>	Primer AP2 for PCR using cDNA of Arabidopsis thaliana e Columbia containing adapter sequences ligated to both i ends	
	<400>	5	
	actcactata (gggetegage gge	23
	<210>	6	
IJ		30	
en de	<211>	DNA	
Fil			
the free from half from mail form	<213>	Artificial sequence	
	<220>		
Ħ,	<223>	MSH3 specific primer S525 for PCR using cDNA of Arabido	nsis
	1200	thaliana ecotype Columbia	70210
164 144 144	<400>	6	
Mindre Company	aggttctgat	tatgtgtgac gctttactta	30
	<210>	7	
	<211>	29	
	<212>		
		DNA Denificial common as	
	<213>	Artificial sequence	
	<220>		
	<223>	MSH3 specific primer S51 for PCR using cDNA of Arabidor thaliana ecotype Columbia	sis
,	<400>	7	
	ggatcgggta	ctgggttttg agtgtgagg	29
ŧ	0.00		
	<210>	8	
	<211>	24	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
		MCIIO anonifia maima- con se man	
	<223>	MSH3 specific primer 635 for PCR using cDNA of Arabidop thaliana ecotype Columbia	psis

```
8
<400>
                                                                      24
gcacgtgctt gatggtgttt tcac
<210>
             9
<211>
            28
            DNA
<212>
             Artificial sequence
<213>
<220>
             MSH3 specific primer S523 for PCR using cDNA of Arabidopsis
<223>
             thaliana ecotype Columbia
<400>
                                                                      28
tcagacagta tccagcatgg cagaagta
<210>
             10
<211>
             33
            DNA
<21.2>
             Artificial sequence
<213>
<220>
             MSH3 specific primer 1S5 for PCR using cDNA of Arabidopsis
<223>
             thaliana ecotype Columbia
             10
<400>
                                                                       33
atcccgggat gggcaagcaa aagcagcaga cga
<210>
             11
             27
<211>
<212>
            DNA
<213>
             Artificial sequence
<220>
             MSH3 specific primer S53 for PCR using cDNA of Arabidopsis
<223>
             thaliana ecotype Columbia
<400>
             11
                                                                       27
gacaaagagc gaaatgaggc cccttgg
<210>
              12
             1250
<211>
<212>
             Arabidopsis thaliana ecotype Columbia
 <213>
<223>
             Clone 52
```

1.1 1997 1911

1

PCT/EP98/06977

12

<400>

cccgggatgg	gcaagcaaaa	gcagcagacg	atttctcgtt	tcttcgctcc	caaacccaaa	60
tccccgactc	acgaaccgaa	tccggtagcc	gaatcatcaa	caccgccacc	gaagatatcc	120
gccactgtat	ccttctctcc	ttccaagcgt	aagcttctct	ccgaccacct	cgccgccgcg	180
tcacccaaaa	agcctaaact	ttctcctcac	actcaaaacc	cagtacccga	tcccaattta	240
caccaaagat	ttctccagag	atttctggaa	ccctcgccgg	aggaatatgt	tcccgaaacg	300
tcatcatcga	ggaaatacac	accattggaa	cagcaagtgg	tggagctaaa	gagcaagtac	360
ccagatgtgg	ttttgatggt	ggaagttggt	tacaggtaca	gattcttcgg	agaagacgcg	420
gagatcgcag	cacgcgtgtt	gggtatttac	gctcatatgg	atcacaattt	catgacggcg	480
agtgtgccaa	catttcgatt	gaatttccat	gtgagaagac	tggtgaatgc	aggatacaag	540
attggtgtag	tgaagcagac	tgaaactgca	gccattaagt	cccatggtgc	aaaccggacc	600
ggcccttttt	teeggggaet	gtcggcgttg	tataccaaag	ccacgcttga	agcggctgag	660
gatataagtg	gtggttgtgg	tggtgaagaa	ggttttggtt	cacagagtaa	tttcttggtt	720
tgtgttgtgg	atgagagagt	taagtcggag	acattaggct	gtggtattga	aatgagtttt	780
gatgttagag	teggtgttgt	tggcgttgaa	atttcgacag	gtgaagttgt	ttatgaagag	840
ttcaatgata	atttcatgag	aagtggatta	gaggctgtga	ttttgagctt	gtcaccagct	900
gagetgttge	ttggccagcc	tctttcacaa	caaactgaga	agtttttggt	ggcacatgct	960
ggacctacct	caaacgttcg	agtggaacgt	gcctcactgg	attgtttcag	caatggtaat	1020
gcagtagatg	aggttatttc	attatgtgaa	aaaatcagcg	caggtaactt	agaagatgat	1080
aaagaaatga	agctggaggc	tgctgaaaaa	ggaatgtctt	gcttgacagt	tcatacaatt	1140
atgaacatgc	cacatetgae	tgttcaagcc	ctcgccctaa	cgttttgcca	tctcaaacag	1200
tttggatttg	aaaggatcct	ttaccaaggg	gcctcatttc	gctctttgtc		1250

<210> 13

<211> 34

<21.2> DNA

<213> Artificial sequence

<220>

<223> MSH3 specific primer 2S5 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia

1.1

13

<400>

atcccgggtc (aaaatgaaca a	agttggtttt	agtc			34
<210>	14					
<211>	27					
<212>	DNA					
<213>	Artificia:	l sequence				
<220>						
<223>		ific primer ecotype Col		R using cDN	A of Arabido	psis
<400>	14					
gccacatctg	actgttcaag	ccctcgc				27
<210>	15					
<211>	2110					
<212>	DNA			7		
	Arabidops	is thailana	ecotype co	Tumbla		
<223>	Clone 13					
<400>	15					
gccacatctg	actgttcaag	ccctcgccct	aacgttttgc	catctcaaac	agtttggatt	60
tgaaaggatc	ctttaccaag	gggcctcatt	tcgctctttg	tcaagtaaca	cagagatgac	120
tctctcagcc	aatactctgc	aacagttgga	ggttgtgaaa	aataattcag	atggatcgga	180
atctggctcc	ttattccata	atatgaatca	cacacttaca	gtatatggtt	ccaggettet	240
tagacactgg	gtgactcatc	ctctatgcga	tagaaatttg	atatctgctc	ggcttgatgc	300
tgtttctgag	atttctgctt	gcatgggatc	tcatagttct	teccagetca	gcagtgagtt	360
ggttgaagaa	ggttctgaga	gagcaattgt	atcacctgag	ttttatctcg	tgctctcctc	420
agtcttgaca	gctatgtcta	gatcatctga	tattcaacgt	ggaataacaa	gaatctttca	480
teggaetget	aaagccacag	agttcattgc	agttatggaa	gctattttac	ttgcggggaa	540
gcaaattcag	cggcttggca	taaagcaaga	ctctgaaatg	aggagtatgc	aatctgcaac	600
tgtgcgatct	actcttttga	gaaaattgat	ttctgttatt	tcatcccctg	ttgtggttga	660
caatgccgga	aaacttctct	ctgccctaaa	taaggaagcg	gctgttcgag	gtgacttgct	720
cgacatacta	atcacttcca	gcgaccaatt	tcctgagctt	gctgaagctc	gccaagcagt	780
tttagtcatc	agggaaaagc	tggattcctc	gatagettea	tttcgcaaga	agctcgctat	840

The second of th

The state of the s

WO 99/19492 PCT/EP98/06977

7

tcgaaatttg	gaatttcttc	aagtgtcggg	gatcacacat	ttgatagagc	tgcccgttga	900
ttccaaggtc	cctatgaatt	gggtgaaagt	aaatagcacc	aagaagacta	ttcgatatca	960
teccecagaa	atagtagctg	gcttggatga	gctagctcta	gcaactgaac	atcttgccat	1020
tgtgaaccga	gcttcgtggg	atagtttcct	caagagtttc	agtagatact	acacagattt	1080
taaggctgcc	gttcaagctc	ttgctgcact	ggactgtttg	cactcccttt	caactctatc	1140
tagaaacaag	aactatgtcc	gtcccgagtt	tgtggatgac	tgtgaaccag	ttgagataaa	1200
catacagtct	ggtcgtcatc	ctgtactgga	gactatatta	caagataact	tcgtcccaaa	1260
tgacacaatt	ttgcatgcag	aaggggaata	ttgccaaatt	atcaccggac	ctaacatggg	1320
aggaaagagc	tgctatatcc	gtcaagttgc	tttaatttcc	ataatggctc	aggttggttc	1380
ctttgtacca	gcgtcattcg	ccaagctgca	cgtgcttgat	ggtgttttca	ctcggatggg	1440
tgcttcagac	agtatccagc	atggcagaag	tacctttcta	gaagaattaa	gtgaagcgtc	1500
acacataatc	agaacctgtt	cttctcgttc	gcttgttata	ttagatgagc	ttggaagagg	1560
cactagcaca	cacgacggtg	tagccattgc	ctatgcaaca	ttacagcatc	tcctagcaga	1620
aaagagatgt	ttggttcttt	ttgtcacgca	ttaccctgaa	atagctgaga	tcagtaacgg	1680
attcccaggt	tctgttggga	cataccatgt	ctcgtatctg	acattgcaga	aggataaagg	1740
cagttatgat	catgatgatg	tgacctacct	atataagctt	gtgcgtggtc	tttgcagcag	1800
gagctttggt	tttaaggttg	ctcagcttgc	ccagatacct	ccatcatgta	tacgtcgagc	1860
catttcaatg	gctgcaaaat	tggaagctga	ggtacgtgca	agagagagaa	atacacgcat	1920
gggagaacca	gaaggacatg	aagaaccgag	aggcgcagaa	gaatctattt	cggctctagg	1980
tgacttgttt	gcagacctga	aatttgctct	ctctgaagag	gacccttgga	aagcattcga	2040
gtttttaaag	catgcttgga	agattgctgg	caaaatcaga	ctaaaaccaa	cttgttcatt	2100
ttgacccggg						2110

<2±0>	16
<211>	29
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer S51 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia
<400>	16

and the second of the second o

	ggato	gggt	ta ct	ggg	tttt	g agt	gtga	agg									29
.	<210 s <211 s <212 s <213 s	> >		17 30 DNA Art:	ific:	ial s	seque	ence									
*	<220: <223:				3 spe liana						or PO	CR us	sing	CDN	A of	Arabid	opsis
	<400:	>		17													
	aggt	tctg	at t	atgt	gtga	c gc	ttta	ctta									30
the state of the s	<210 <211 <212 <213 <220 <221 <222 <223	> > > > > > > > > > > > > > > > > > > >		CDS (10 AtM pol	bido 0)	(3 full	342) -len							uenc	e of	the er	ncoded
	<400			18											12.00	octtc	60
de compa dell'accomp dell'acco												cage	Cace	jac c	.acge	iccttc	99
					ggagg												
	atg Met 1	Gly	aag Lys	caa Gln	aag Lys 5	cag Gln	cag Gln	acg Thr	att Ile	Ser 10	cgt Arg	Phe	Phe	gct Ala	Pro 15	aaa Lys	147
					act Thr												195
•					ata Ile												243
ir			Leu		gac Asp												291
					act Thr		Asn					Pro					339

 $(a_{i,j},a_{$

													gtt Val 95		387
	-	_			-								gtg Val		435
			-	-	-								gtt Val		483
													cgc Arg		531
													agt Ser		579
				_	-					_	_		gca Ala 175		627
y Sign													aag Lys		675
													gcg Ala		723
													ggt Gly		771
							Gly				Phe		tgt Cys		819
•	_				-	Lys				Gly			gaa Glu 255		867
æ,	_			_		_		_	Gly	_	_			ggt Gly	915
	-			Tyr	-	-		Asp			_	Ser		tta Leu	963

and the control of the complete many control of the
	gag Glu	gct Ala 290	gtg Val	att Ile	ttg Leu	agc Ser	ttg Leu 295	tca Ser	cca Pro	gct Ala	gag Glu	ctg Leu 300	ttg Leu	ctt Leu	ggc Gly	cag Gln	1011
•															gga Gly		1059
*	acc Thr	tca Ser	aac Asn	gtt Val	cga Arg 325	gtg Val	gaa Glu	cgt Arg	gcc Ala	tca Ser 330	ctg Leu	gat Asp	tgt Cys	ttc. Phe	agc Ser 335	aat Asn	1107
															agc Ser		1155
Section 2															gaa Glu		1203
County daily, may, sing															cat His		1251
												Leu			ttt Phe		1299
											Phe				tca Ser 415		1347
					Thr					Thr					gag Glu		1395
				Asn					Glu					. Phe	cat His		1443
÷			His					Tyr					Leu		cac His		1491
₽		Thr				-	Asp					e Ser	_		ctt Leu		1539
	_	_				Ser	-		-	-	seı Seı					cag Gln	1587

								1 (
							ggt Gly 505								1635
							tca Ser								1683
							aca Thr								1731
Ly							atg Met								1779
							aag Lys								1827
							act Thr 585								1875
_							gac Asp								1923
							cga Arg								1971
I							gag Glu			Glu					2019
							gat Asp		Ser					Arg	2067
				Ile			gaa Glu 665	Phe					Gly	atc Ile	2115
			ı Ile				. Asp					His		tgg Trp	2163
		. Val				Lys					: His			gaa Glu	2211

ata gta Ile Val 705	gct gg Ala Gl	y Leu	gat g Asp G 710	gag o Glu I	cta (Leu .	gct (Ala)	Leu	gca Ala 715	act Thr	gaa Glu i	cat His	Leu	gcc Ala 720	2259
att gtg Ile Val	aac cg Asn Ar	a gct g Ala 725	tcg t Ser T	gg g Irp A	gat Asp	Ser	ttc Phe 730	ctc Leu	aag Lys	agt Ser	ttc Phe	agt Ser 735	aga Arg	2307
tac tac Tyr Tyr		p Phe			Ala									2355
tgt ttg Cys Leu	cac tc His Se 755	c ctt r Leu	tca a	Thr :	cta Leu 760	tct Ser	aga Arg	aac Asn	aag Lys	aac Asn 765	tat Tyr	gtc Val	cgt Arg	2403
ccc gag Pro Glu 770	ttt gt Phe Va	g gat	Asp (tgt Cys 775	gaa Glu	cca Pro	gtt Val	gag Glu	ata Ile 780	aac Asn	ata Ile	cag Gln	tct Ser	2451
ggt cgt Gly Arg 785	cat co	et gta co Val	ctg (Leu (gag Glu	act Thr	ata Ile	tta Leu	caa Gln 795	gat Asp	aac Asn	ttc Phe	gtc Val	cca Pro 800	2499
aat gac Asn Asp														2547
gga cct Gly Pro	aac at Asn Me 82	et Gly	gga Gly	aag Lys	agc Ser	tgc Cys 825	tat Tyr	atc Ile	cgt Arg	caa Gln	gtt Val 830	gct Ala	tta Leu	2595
att tcc Ile Ser	ata at Ile Me 835	tg gct et Ala	cag Gln	gtt Val	ggt Gly 840	tcc Ser	ttt Phe	gta Val	cca Pro	gcg Ala 845	tca Ser	ttc Phe	gcc Ala	2643
aag ctg Lys Leu 850	His Va									Gly				2691
agt ato Ser Ile 865									Glu					2739
tca cac Ser His			Thr					Ser					Asp	2787
gag ctt Glu Lev	ı Gly A	ga ggo rg Gly	act Thr	agc Ser	aca Thr	cac His 905	Asp	ggt Gly	gta Val	gcc Ala	att Ile 910	Ala	tat Tyr	2835

can find the control of the

	gca Ala	aca Thr	tta Leu 915	cag Gln	cat His	ctc Leu	cta Leu	gca Ala 920	gaa Glu	aag Lys	aga Arg	tgt Cys	ttg Leu 925	gtt Val	ctt Leu	ttt Phe	2883
4	gtc Val	acg Thr 930	cat His	tac Tyr	cct Pro	gaa Glu	ata Ile 935	gct Ala	gag Glu	atc Ile	agt Ser	aac Asn 940	gga Gly	ttc Phe	cca Pro	ggt Gly	2931
.	tct Ser 945	gtt Val	g1y 999	aca Thr	tac Tyr	cat His 950	gtc Val	tcg Ser	tat Tyr	ctg Leu	aca Thr 955	ttg Leu	cag Gln	aag Lys	gat Asp	aaa Lys 960	2979
	ggc Gly	agt Ser	tat Tyr	gat Asp	cat His 965	gat Asp	gat Asp	gtg Val	acc Thr	tac Tyr 970	cta Leu	tat Tyr	aag Lys	ctt Leu	gtg Val 975	cgt Arg	3027
	ggt Gly	ctt Leu	tgc Cys	agc Ser 980	agg Arg	agc Ser	ttt Phe	ggt Gly	ttt Phe 985	aag Lys	gtt Val	gct Ala	cag Gln	ctt Leu 990	gcc Ala	cag Gln	3075
	ata Ile	cct Pro	cca Pro 995	tca Ser	tgt Cys	ata Ile	Arg	cga Arg 1000	Ala	att Ile	tca Ser	Met	gct Ala 1005	gca Ala	aaa Lys	ttg Leu	3123
	Glu	gct Ala 1010	gag Glu	gta Val	cgt Arg	Ala	aga Arg 1015	Glu	aga Arg	aat Asn	aca Thr	cgc Arg 1020	atg Met	gga Gly	gaa Glu	cca Pro	3171
	gaa Glu 102	Gly	cat His	gaa Glu	Glu	ccg Pro 1030	Arg	ggc	gca Ala	gaa Glu	gaa Glu 1035	Ser	att Ile	tcg Ser	gct Ala	cta Leu 1040	3219
	ggt Gly	gac Asp	ttg Leu	ttt Phe	gca Ala 1045	Asp	ctg Leu	aaa Lys	ttt Phe	gct Ala	Leu	tct Ser	gaa Glu	gag Glu	gac Asp	Pro	3267
	tgg Trp	aaa Lys	gca Ala	ttc Phe	Glu	ttt Phe	tta Leu	aag Lys	cat His 1065	Ala	tgg Trp	g aag Lys	g att	gct Ala 1070	ggc	aaa Lys	3315
ı.				Lys					r Phe		attta	aatc	ttaa	acatt	at		3362
	ago	aact	gca	aggt	cttg	gat o	catct	gtta	ag tt	gegi	tacta	a act	tate	gtgt	atta	agtataa	3422
9	caa	ıgaaa	aga	gaat	taga	aga g	gatgo	gatto	ct aa	atcc	ggtgi	t tg	cagta	acat	cttt	tctcca	3482
	cco	gcat	aaa	aaaa	aaaa	aaa a	aaaa	aaaa	aa aa	aaaa	aaaa	a					3522
		.0.		7 0	,												

The second secon

<210> 19 <211> 1081 <212> PRT

 $(r_{i+1}, \ldots, r_{i+1}, \ldots, r_{$

	<213 <223				bido ypep				a ec	otyp	e Co	lumb	ia			
	<400	>		19												
	Met 1	Gly	Lys	Gln	Lys 5	Gln	Gln	Thr	Ile	Ser 10	Arg	Phe	Phe	Ala	Pro 15	Lys
	Pro	Lys	Ser	Pro 20	Thr	His	Glu	Pro	Asn 25	Pro -	Val	Ala	Glu	Ser 30	Ser	Thr
	Pro	Pro	Pro 35	Lys	Ile	Ser	Ala	Thr 40	Val	Ser	Phe	Ser	Pro 45	Ser	Lys	Arg
	Lys	Leu 50	Leu	Ser	Asp	His	Leu 55	Ala	Ala	Ala	Ser	Pro 60	Lys	Lys	Pro	Lys
	Leu 65	Ser	Pro	His	Thr	Gln 70	Asn	Pro	Val	Pro	Asp 75	Pro	Asn	Leu	His	Gln 80
	Arg	Phe	Leu	Gln	Arg 85	Phe	Leu	Glu	Pro	Ser 90	Pro	Glu	Glu	Tyr	Val 95	Pro
The state of the s	Glu	Thr	Ser	Ser 100	Ser	Arg	Lys	Tyr	Thr 105	Pro	Leu	Glu	Gln	Gln 110	Val	Val
	Glu	Leu	. Lys 115		Lys	Tyr	Pro	Asp 120	Val	Val	Leu	Met	Val 125	Glu	Val	Gly
1000	Tyr	Arg		Arg	Phe	Phe	Gly 135		Asp	Ala	. Glu	Ile 140		Ala	Arg	Val
	Leu 145		/ Ile	Tyr	Ala	His 150		Asp	His	Asr	Phe		Thr	· Ala	Ser	Val 160
	Pro) Thr	Phe	e Arg	Leu 165		Phe	His	Val	Arg		Leu	. Val	. Asn	. Ala 175	Gly
	Туг	: Lys	s Ile	e Gly 180		Val	Lys	Gln	Thr 185		ı Thr	Ala	Ala	190		Ser
۵	His	; Gly	y Ala 195		Arg	Thr	: Gly	/ Pro 200		Phe	e Arg	gly	/ Let 205		Ala	. Leu
5	Туз	r Th:		s Ala	Thr	Let	1 Glu 215		a Ala	a Gli	ı Asp	220		c Gly	/ Gly	/ Cys
	Gl ₃ 22		y Gl	u Glu	ı Gly	/ Phe		y Sei	r Gli	ı Se	r Ası 239		e Le	ı Val	L Cys	3 Val 240
	Va	l As	p Gl	u Arg	y Val 245		s Sei	r Gli	u Thi	r Le		y Cys	s Gl	y Ile	e Glu 25	ı Met 5

The state of the s

WO 99/19492 PCT/EP98/06977

15

Ser	Phe	Asp	Val 260	Arg	Val	Gly	Val	Val 265	Gly	Val	Glu	Ile	Ser 270	Thr	Gly
Glu	Val	Val 275	Tyr	Glu	Glu	Phe	Asn 280	qaA	Asn	Phe	Met	Arg 285	Ser	Gly	Leu
Glu	Ala 290	Val	Ile	Leu	Ser	Leu 295	Ser	Pro	Ala	Glu	Leu 300	Leu	Leu	Gly	Gln
Pro 305	Leu	Ser	Gln	Gln	Thr 310	Glu	Lys	Phe	Leu	Val 315	Ala	Met	Ala	Gly	Pro 320
Thr	Ser	Asn	Val	Arg 325	Val	Glu	Arg	Ala	Ser 330	Leu	Asp	Cys	Phe	Ser 335	Asn
Gly	Asn	Ala	Val 340	Asp	Glu	Val	Ile	Ser 345	Leu	Cys	Glu	Lys	Ile 350	Ser	Ala
Gly	Asn	Leu 355	Glu	Asp	Asp	Lys	Glu 360	Met	Lys	Leu	Glu	Ala 365	Ala	Glu	Lys
Gly	Met 370	Ser	Cys	Leu	Thr	Val 375	His	Thr	Ile	Met	Asn 380	Met	Pro	His	Leu
Thr 385	Val	Gln	Ala	Leu	Ala 390	Leu	Thr	Phe	Cys	His 395	Leu	Lys	Gln	Phe	Gly 400
Phe	Glu	Arg	Ile	Leu 405	Tyr	Gln	Gly	Ala	Ser 410	Phe	Arg	Ser	Leu	Ser 415	Ser
Asn	Thr	Glu	Met 420	Thr	Leu	Ser	Ala	Asn 425	Thr	Leu	Gln	Gln	Leu 430	Glu	Val
Val	Lys	Asn 435	Asn	Ser	Asp	Gly	Ser 440	Glu	Ser	Gly	Ser	Leu 445	Phe	His	Asn
Met	Asn 450	His	Thr	Leu	Thr	Val 455	_	Gly	Ser	Arg	Leu 460		Arg	His	Trp
Val 465	Thr	His	Pro	Leu	Cys 470	Asp	Arg	Asn	Leu	Ile 475	Ser	Ala	Arg	Leu	Asp 480
Ala	Val	Ser	Glu	Ile 485	Ser	Ala	Cys	Met	Gly 490		His	Ser	Ser	Ser 495	Gln
Leu	Ser	Ser	Glu 500		Val	Glu	Glu	Gly 505		Glu	Arg	Ala	Ile 510	Val	Ser
Pro	Glu	Phe 515	_	Leu	Val	Leu	Ser 520		Val	. Leu	Thr	Ala 525	Met	Ser	Arg
Ser	Ser 530	_	Ile	Gln	Arg	Gly 535		Thr	Arg	, Ile	Phe		Arg	Thr	Ala

the state of the s

and the second of the second o

	Lys 545	Ala	Thr	Glu	Phe	Ile 550	Ala	Val	Met	Glu	Ala 555	Ile	Leu	Leu	Ala	Gly 560
	Lys	Gln	Ile	Gln	Arg 565	Leu	Gly	Ile	Lys	Gln 570	Asp	Ser	Glu	Met	Arg 575	Ser
*	Met	Gln	Ser	Ala 580	Thr	Val	Arg	Ser	Thr 585	Leu	Leu	Arg	Lys	Leu 590	Ile	Ser
	Val	Ile	Ser 595	Ser	Pro	Val	Val	Val 600	Asp	Asn	Ala	Gly	Lys 605	Leu	Leu	Ser
	Ala	Leu 610	Asn	Lys	Glu	Ala	Ala 615	Val	Arg	Gly	Asp	Leu 620	Leu	Asp	Ile	Leu
	Ile 625	Thr	Ser	Ser	Asp	Gln 630	Phe	Pro	Glu	Leu	Ala 635	Glu	Ala	Arg	Gln	Ala 640
	Val	Leu	Val	Ile	Arg 645	Glu	Lys	Leu	Asp	Ser 650	Ser	Ile	Ala	Ser	Phe 655	Arg
	Lys	Lys	Leu	Ala 660	Ile	Arg	Asn	Leu	Glu 665	Phe	Leu	Gln	Val	Ser 670	Gly	Ile
	Thr	His	Leu 675	Ile	Glu	Leu	Pro	Val 680	Asp	Ser	Lys	Val	Pro 685	His	Asn	Trp
	Val	Lys 690	Val	Asn	Ser	Thr	Lys 695	Lys	Thr	Ile	Arg	Tyr 700	His	Pro	Pro	Glu
Maria de la companya	Ile 705	Val	Ala	Gly	Leu	Asp 710		Leu	Ala	Leu	Ala 715	Thr	Glu	His	Leu	Ala 720
	Ile	Val	Asn	Arg	Ala 725		Trp	qsA	Ser	730		Lys	Ser	Phe	Ser 735	Arg
	Tyr	Tyr	Thr	Asp 740		Lys	Ala		Val 745		ı Ala	Leu	Ala	750		Asp
	Cys	Leu	His 755	Ser	Leu	. Ser	Thr	Leu 760		: Arg	y Asn	Lys	765		Val	Arg
°	Pro	Glu 770	Phe	Val	Asp	Asp	Cys 775		Pro	Va]	l Glu	780		ı Ile	Gln	Ser
ð	Gly 785	_	His	Pro	Val	1 Leu 790		Thr	: Ile	e Lei	ı Gln 795) Asr	n Phe	· Val	Pro 800
	Asn	Asp	Thr	Ile	Let 805		: Ala	Glu	ı Gly	/ Gl: 81		Cys	Glr	n Ile	815	Thr
	Gly	Pro	Asn	Met 820		/ Gly	, The	Ser	Cys 825		r Ile	Arç	g Glr	Nal 830		Leu

Ile Ser Ile Met Ala Gln Val Gly Ser Phe Val Pro Ala Ser Phe Ala 835 840 845

Lys Leu His Val Leu Asp Gly Val Phe Thr Arg Met Gly Ala Ser Asp 850 855 860

Ser Ile Gln His Gly Arg Ser Thr Phe Leu Glu Glu Leu Ser Glu Ala 865 870 875 880

Ser His Ile Ile Arg Thr Cys Ser Ser Arg Ser Leu Val Ile Leu Asp 885 890 895

Glu Leu Gly Arg Gly Thr Ser Thr His Asp Gly Val Ala Ile Ala Tyr 900 905 910

Ala Thr Leu Gln His Leu Leu Ala Glu Lys Arg Cys Leu Val Leu Phe 915 920 925

Val Thr His Tyr Pro Glu Ile Ala Glu Ile Ser Asn Gly Phe Pro Gly 930 935 940

Ser Val Gly Thr Tyr His Val Ser Tyr Leu Thr Leu Gln Lys Asp Lys 945 950 955 960

Gly Ser Tyr Asp His Asp Asp Val Thr Tyr Leu Tyr Lys Leu Val Arg 965 970 975

Gly Leu Cys Ser Arg Ser Phe Gly Phe Lys Val Ala Gln Leu Ala Gln 980 985 990

Ile Pro Pro Ser Cys Ile Arg Arg Ala Ile Ser Met Ala Ala Lys Leu
995 1000 1005

Glu Ala Glu Val Arg Ala Arg Glu Arg Asn Thr Arg Met Gly Glu Pro 1010 1015 1020

Glu Gly His Glu Glu Pro Arg Gly Ala Glu Glu Ser Ile Ser Ala Leu 1025 1030 1035 1040

Gly Asp Leu Phe Ala Asp Leu Lys Phe Ala Leu Ser Glu Glu Asp Pro 1045 1050 1055

the state of the company production of the state of the s

Trp Lys Ala Phe Glu Phe Leu Lys His Ala Trp Lys Ile Ala Gly Lys
1060 1065 1070

Ile Arg Leu Lys Pro Thr Cys Ser Phe 1075 1080

<210> 20 <211> 24

<212> DNA

<213> Artificial sequence

		18	_
	<220>	and a partition and Arabidonese	
	<223>	MSH6 specific primer 638 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
	<400>	20	
*	tetetaceag (grgacgaaaa accg	24
÷			
••	<210>	21	
	<211>	28	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Primer S81 for PCR using cDNA of Arabidopsis thaliana ecot Columbia	уре
	<400>	21	
	cgtcgccttt	agcatccct tccttcac	28
W	<210>	22	
	<211>	30	
ā		DNA	
	<212>		
500	<213>	Artificial sequence	
-1	<220>	and a part of Arabidons	
Francisco	<223>	MSH6 specific primer S823 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	.5
	<400>	22	
	gcttggcgca	tctaatagaa tcatgacagg	30
	<210>	23	
	<211>	24	
	<211>	DNA	
	<213>	Artificial sequence	
-	<220>		
•	<223>	MSH6 specific primer 637 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	s
ð	<400>	23	
	gacagcgtca	gttcttcaga atgc	24
	<210>	24	
	<211>	33	
	<212>	DNA	

the control of the complete section of the control
	21.	Numi for mini componed	
	<213>	Artificial sequence	
	<220>		
	<223>	MSH6 specific primer 1S8 for PCR using cDNA of Arabidopsisthaliana ecotype Columbia	S
	<400>	24	
	atcccgggat	gcagcgccag agatcgattt tgt	33
	<210>	25	
	<211>	27	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		_
	<223>	MSH6 specific primer S83 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	S
	<400>	25	
Topic	cgctatctat	ggetgetteg aattgag	27
dla	<210>	26	
	<211>	1385	
	<212>	DNA	
	<213>	Arabidopsis thaliana ecotype Columbia	
74 74	<223>	Clone 43	
in the second se	(22)	0.20.10	
	<400>	26	
	cccgggatgc	agegeeagag ategattitg tetttettee aaaaaceeac ggeggegaet	60
	acgaagggtt	tggtttccgg cgatgctgct agcggcgggg gcggcagcgg aggaccacga	120
	tttaatgtga	aggaagggga tgctaaaggc gacgcttctg tacgttttgc tgtttcgaaa	180
	tctgtcgatg	aggttagagg aacggatact ccaccggaga aggttccgcg tcgtgtcctg	240
	ccgtctggat	ttaagccggc tgaatccgcc ggtgatgett cgtccctgtt ctccaatatt	300
	atgcataagt	ttgtaaaagt cgatgatcga gattgttctg gagagaggag ccgagaagat	360
ŕ	gttgttccgc	tgaatgattc atctctatgt atgaaggcta atgatgttat tcctcaattt	420
	cgttccaata	atggtaaaac tcaagaaaga aaccatgctt ttagtttcag tgggagagct	480
	gaacttagat	cagtagaaga tataggagta gatggcgatg ttcctggtcc agaaacacca	540
	gggatgcgtc	cacgtgcttc tcgcttgaag cgagttctgg aggatgaaat gacttttaag	600
	gaggataagg	ttcctgtatt ggactctaac aaaaggctga aaatgctcca ggatccggtt	660

tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	aatggcttga	gtcttctcga	720
atcagggatg	ccaatagaag	acgtcctgat	gatccccttt	acgatagaaa	gaccttacac	780
ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	agcaatattg	gagtgttaag	840
agtgaatata	tggacattgt	gcttttcttt	aaagtgggga	aattttatga	gctgtatgag	900
ctagatgcgg	aattaggtca	caaggagctt	gactggaaga	tgaccatgag	tggtgtggga	960
aaatgcagac	aggttggtat	ctctgaaagt	gggatagatg	aggcagtgca	aaagctatta	1020
gctcgtggat	ataaagttgg	acgaatcgag	cagctagaaa	catctgacca	agcaaaagcc	1080
agaggtgcta	atactataat	tccaaggaag	ctagttcagg	tattaactcc	atcaacagca	1140
agcgagggaa	acategggee	tgatgccgtc	catcttcttg	ctataaaaga	gatcaaaatg	1200
gagctacaaa	agtgttcaac	tgtgtatgga	tttgcttttg	ttgactgtgc	tgccttgagg	1260
ttttgggttg	ggtccatcag	cgatgatgca	tcatgtgctg	ctcttggagc	gttattgatg	1320
caggtttctc	caaaggaagt	gttatatgac	agtaaagggc	tatcaagaga	agcacaaaag	1380
gctctaagga	aatatacgtt	gacagggtct	acggcggtac	agttggctcc	agtaccacaa	1440
gtaatggggg	atacagatgc	tgctggagtt	agaaatataa	tagaatctaa	. cggatacttt	1500
aaaggttctt	ctgaatcatg	gaactgtgct	gttgatggtc	taaatgaatg	tgatgttgcc	1560
cttagtgctc	ttggagagct	aattaatcat	ctgtctaggc	taaagctaga	agatgtactt	1620
aagcatgggg	atatttttcc	ataccaagtt	tacaggggtt	gtctcagaat	: tgatggccag	1680
acgatggtaa	atcttgagat	atttaacaat	agctgtgatg	gtggtccttc	agggaccttg	1740
tacaaatato	ttgataactg	, tgttagtcca	actggtaagc	gactcttaag	g gaattggatc	1800
tgccatccac	: tcaaagatgt	agaaagcato	c aataaacggc	: ttgatgtagt	tgaagaattc	1860
acggcaaact	: cagaaagtat	gcaaatcact	ggccagtato	: tccacaaact	tccagactta	1920
gaaagactgo	tcggacgcat	caagtctago	c gttcgatcat	cageetetgt	gttgcctgct	1980
cttctgggga	a aaaaagtgc	gaaacaacga	a gttaaagcat	ttgggcaaa	t tgtgaaaggg	2040
ttcagaagto	g gaattgatc	t gttgttggci	t ctacagaagg	g aatcaaata	t gatgagtttg	2100
ctttataaa	tctgtaaac	t tootatatta	a gtaggaaaaa	a gegggetag	a gttatttctt	2160
tctcaattc	g aagcagcca	t agatagcg				2188

The state of the s

PCT/EP98/06977

27

<210>

1385 <211> DNA <212> Arabidopsis thaliana ecotype Columbia <213> <223> Clone 62 <400> 27 60 catcagecte tgtgttgeet getettetgg ggaaaaaagt getgaaacaa egagttaaag 120 catttgggca aattgtgaaa gggttcagaa gtggaattga tctgttgttg gctctacaga 180 aggaatcaaa tatgatgagt ttgctttata aactctgtaa acttcctata ttagtaggaa 240 aaagcgggct agagttattt ctttctcaat tcgaagcagc catagatagc gactttccaa attatcagaa ccaagatgtg acagatgaaa acgctgaaac tctcacaata cttatcgaac 300 tttttatcga aagagcaact caatggtctg aggtcattca caccataagc tgcctagatg 360 420 tectgagate ttttgcaate geageaagte tetetgetgg aageatggee aggeetgtta 480 tttttcccga atcagaagct acagatcaga atcagaaaac aaaagggcca atacttaaaa 540 tccaaggact atggcatcca tttgcagttg cagccgatgg tcaattgcct gttccgaatg 600 atatactcct tggcgaggct agaagaagca gtggcagcat tcatcctcgg tcattgttac 660 tgacgggacc aaacatgggc ggaaaatcaa ctcttcttcg tgcaacatgt ctggccgtta 720 tetttgecca acttggetge taegtgeegt gtgagtettg egaaatetee etegtggata 780 ctatcttcac aaggettgge geatetgata gaateatgae aggagagagt acetttttgg 840 tagaatgcac tgagacageg teagttette agaatgcaae teaggattea etagtaatee 900 ttqacqaact qggcaqagga actagtactt tcgatggata cgccattgca tactcggttt ttegteacet ggtagagaaa gtteaatgte ggatgetett tgeaacacat taccaccete 960 tcaccaagga attcgcgtct cacccacgtg tcacctcgaa acacatggct tgcgcattca 1020 1080 aatcaagatc tgattatcaa ccacgtggtt gtgatcaaga cctagtgttc ttgtaccgtt 1140 taaccgaggg agcttgtcct gagagctacg gacttcaagt ggcactcatg gctggaatac 1200 caaaccaagt ggttgaaaca gcatcaggtg ctgctcaagc catgaagaga tcaattgggg aaaacttcaa gtcaagtgag ctaagatctg agttctcaag tctgcatgaa gactggctca 1260 1320 agtcattggt gggtatttct cgagtcgccc acaacaatgc ccccattggc gaagatgact 1380 acgacacttt gttttgctta tggcatgaga tcaaatcctc ttactgtgtt cccaaataac

1.11

The state of the s

PCT/EP98/06977

	cc g gg	138	5
	<210>	28	
	<211>	34	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	MSH6 specific primer 2S8 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
	<400>	26	
	atcccgggtt	atttgggaac acagtaagag gatt	34
ì			
1	<210>	29	
F	<211>	27	
: :	<212>	DNA	
	<213>	Artificial sequence	
ļ 1	<220>		
	<223>	MSH6 specific primer S82 for PCR using cDNA of Arabidopsis	
	, , , , , , , , , , , , , , , , , , , ,	thaliana ecotype Columbia	
	<400>	29	
M 40015- Minut 3111	gcgttcgatc	atcagcctct gtgttgc	27
	<210>	30	
	<211>	3606	
	<212>	DNA	
	<213>	Arabidopsis thaliana ecotype Columbia	
	<220>		
	<221>	CDS	
	<222>	(142)(3468)	
	<223>	AtMSH6 full-length cDNA and deduced sequence of the encoded	4
	(223)	polypeptide	^
	<400>	30	
	aaaagttgag	ccctgaggag tatcgtttcc gccatttcta cgacgcaagg cgaaaatttt 6	0
	tggcgccaat	ctttccccc tttcgaattc tctcagctca aaacatcgtt tctctctcac 12	0
1	tctctctcac	aattccaaaa a atg cag cgc cag aga tcg att ttg tct ttc Met Gln Arg Gln Arg Ser Ile Leu Ser Phe 1 5 10	1

				ccc Pro													219
¥				ggc 30													267
£	gaa Glu	GJÀ aaa	gat Asp 45	gct Ala	aaa Lys	ggc Gly	gac Asp	gct Ala 50	tct Ser	gta Val	cgt Arg	ttt Phe	gct Ala 55	gtt Val	tcg Ser	aaa Lys	315
	tct Ser	gtc Val 60	gat Asp	gag Glu	gtt Val	aga Arg	gga Gly 65	acg Thr	gat Asp	act Thr	cca Pro	ccg Pro 70	gag Glu	aag Lys	gtt Val	ccg Pro	363
				ctg Leu													411
				ctg Leu													459
	gat Asp	cga Arg	gat Asp	tgt Cys 110	tct Ser	gga Gly	gag Glu	agg Arg	agc Ser 115	cga Arg	gaa Glu	gat Asp	gtt Val	gtt Val 120	ccg Pro	ctg Leu	507
STATE OF THE PROPERTY OF THE P				tct Ser										Pro			555
	cgt Arg	tcc Ser 140	Asn	aat Asn	ggt Gly	aaa Lys	act Thr 145	caa Gln	gaa Glu	aga Arg	aac Asn	cat His 150	Ala	ttt Phe	agt Ser	ttc Phe	603
							Arg					Ile				ggc Gly 170	651
K						Glu					. Arg					cgc Arg	699
£					Leu					Thr					Lys	gtt Val	747
				. Asp					j Lev					ı Asp		ggtt Val	795

WO 99/19492 PCT/EP98/06977

24

											aaa Lys 230					843
											cgt Arg					891
											gat Asp					939
											aag Lys					987
											tat Tyr					1035
											tgg Trp 310					1083
											tct Ser					1131
										Gly	tat Tyr				Arg	1179
٠				Glu					Ala		gcc Ala			Ala	aat Asn	1227
			Pro					Gln					Ser		gca Ala	1275
		Gly					Asp					Leu			aaa Lys	1323
	 Ile		_			Gln	_	_			Val				gct Ala 410	1371
					Ala					Val					gat Asp	1419

The second secon

									•	2.5							
	gat Asp	gca Ala	tca Ser	tgt Cys 430	gct Ala	gct Ala	ctt Leu	gga Gly	gcg Ala 435	tta Leu	ttg Leu	atg Met	cag Gln	gtt Val 440	tct Ser	cca Pro	1467
4	aag Lys	gaa Glu	gtg Val 445	tta Leu	tat Tyr	gac Asp	agt Ser	aaa Lys 450	gly ggg	cta Leu	tca Ser	aga Arg	gaa Glu 455	gca Ala	caa Gln	aag Lys	1515
•	gct Ala	cta Leu 460	agg Arg	aaa Lys	tat Tyr	acg Thr	ttg Leu 465	aca Thr	gly aaa	tct Ser	acg Thr	gcg Ala 470	gta Val	cag Gln	ttg Leu	gct Ala	1563
	cca Pro 475	gta Val	cca Pro	caa Gln	gta Val	atg Met 480	gly aaa	gat Asp	aca Thr	gat Asp	gct Ala 485	gct Ala	gga Gly	gtt Val	aga Arg	aat Asn 490	1611
						gga Gly											1659
						cta Leu											1707
	gga Gly	gag Glu	cta Leu 525	att Ile	aat Asn	cat His	ctg Leu	tct Ser 530	agg Arg	cta Leu	aag Lys	cta Leu	gaa Glu 535	gat Asp	gta Val	ctt Leu	1755
			Gly			ttt Phe		Tyr					Gly				1803
	att Ile 555	Asp	ggc Gly	cag Gln	acg Thr	atg Met 560	gta Val	aat Asn	ctt Leu	gag Glu	ata Ile 565	: Phe	aac Asn	aat Asn	agc Ser	tgt Cys 570	1851
						. Gly					Tyr					Val	1899
*					Lys	cga Arg				Ası					Pro		1947
ć				. Glu		ato			arç					l Gli			1995
			a Ası					Gli					а Туз			c aaa E Lys	2043

The state of the s

	-		-	aga Arg 640	_			-		_		_	_	-	 2091
	_		_	ttg Leu		_		_				-	_		2139
				ttt Phe											2187
_	_	_	_	gct Ala		_	_	_			_	-		_	2235
			_	aaa Lys								_			2283
				caa Gln 720		-	-	_		_	_	_			2331
				gat Asp											2379
		-		ttt Phe				-						_	2427
			-	tgc Cys			_	_	_			_		-	2475
				gga Gly											2523
				cag Gln 800											2571
				cat His											2619
				ata Ile											2667

															ggc Gly		2715
						-									gcc Ala		2763
															gtg Val		2811
										-					gga Gly 905		2859
mid find find	_			_	_	_	-								cag Gln		2907
thuit thun thun thuit thun m															gga Gly		2955
=	_			_			_		-		_	_			cac His		3003
of Andrika Mener Term, though	_	-		_		_		_			_				cac His		3051
															cac His 985		3099
	_	-	_				-						Arg		tgt Cys		3147
		Asp					Tyr	-	Leu			Gly	_	Cys		gag Glu	3195
	Ser		Gly			Val	_		-	_	Gly		Pro		caa Gln	gtg Val	3243
	-	Glu			Ser		Ala			Ala		Lys			Ile	999 Gly 1050	3291

WO 99/19492 PCT/EP98/06977

28

	gga Glu	aac Asn	ttc a	Lys :	ca Ser 055	agt (Ser (gag Glu	cta . Leu .	Arg	tct 9 Ser (gag (Glu)	ttc Phe	ca a Ser S	Ser I	ctg Leu 1 065	cat His	3339
	gaa Glu	gac Asp	Trp	ctc a Leu : 070	aag Lys	tca Ser	ttg Leu	Val	ggt Gly 075	att i	ser .	cga (Arg	Val 1	gcc (Ala 1 080	cac . His .	aac Asn	3387
		gcc Ala 1					Asp					Leu					3435
	His	gag Glu 100				Ser						taaa	tggc	ta			3478
	tgac	cataa	ca c	tatc	tgaa	g ct	.cgtt	aagt	ctt	ttgc	ctc	tctg	atgt	tt a	ttcc	tctta	3538
5 10 10 10 10 10 10 10 10 10 10 10 10 10	aaaa	aatgo	tt a	tata	tcaa	ıa aa	atto	rtttc	: ctc	gatt	aaa	aaaa	aaaa	aa a	aaaa	aaaaa	3598
	aaaa	aaaa															3606
	<21: <21: <21: <22: <40:	2 > 3 > 3 >			ubido	opsis otide			ıa ec	cotyp	e Cc	lumb	ia				
	Met	Gln	Arg	Gln	Arg 5	Ser	Ile	Leu	Ser	Phe 10	Phe	Gln	Lys	Pro	Thr 15	Ala	
	Ala	Thr	Thr	Lys 20	Gly	Leu	Val	Ser	Gly 25	Asp	Ala	Ala	Ser	Gly 30	Gly	Gly	
	Gly	Ser	Gly 35	Gly	Pro	Arg	Phe	Asn 40	Val	Arg	Glu	Gly	Asp 45	Ala	Lys	Gly	
	Asp	Ala 50	Ser	Val	Arg	Phe	Ala 55	Val	Ser	Lys	Ser	Val 60	Asp	Glu	Val	Arg	
•	Gly 65	Thr	Asp	Thr	Pro	Pro 70	Glu	Lys	Val	Pro	Arg 75	Arg	Val	Leu	Pro	Ser 80	
	Glv	Phe	Lvs	Pro	Ala	Glu	Ser	Ala	Gly	Asp 90	Ala	Ser	Ser	Leu	Phe 95	Ser	
÷			•		85					90					د ر		
ť		ı Ile				Phe	Val	Lys	Val 105	Asp	Asp	Arg	Asp	Cys 110		Gly	

Met Lys Ala Asn Asp Val Ile Pro Gln Phe Arg Ser Asn Asn Gly Lys 140 130 Thr Gln Glu Arg Asn His Ala Phe Ser Phe Ser Gly Arg Ala Glu Leu 155 150 145 Arg Ser Val Glu Asp Ile Gly Val Asp Gly Asp Val Pro Gly Pro Glu 170 165 Thr Pro Gly Met Arg Pro Arg Ala Ser Arg Leu Lys Arg Val Leu Glu 185 Asp Glu Met Thr Phe Lys Glu Asp Lys Val Pro Val Leu Asp Ser Asn 195 Lys Arg Leu Lys Met Leu Gln Asp Pro Val Cys Gly Glu Lys Lys Glu 215 210 Val Asn Glu Gly Thr Lys Phe Glu Trp Leu Glu Ser Ser Arg Ile Arg 235 230 Asp Ala Asn Arg Arg Pro Asp Asp Pro Leu Tyr Asp Arg Lys Thr 245 Leu His Ile Pro Pro Asp Val Phe Lys Lys Met Ser Ala Ser Gln Lys 260 265 Gln Tyr Trp Ser Val Lys Ser Glu Tyr Met Asp Ile Val Leu Phe Phe Lys Val Gly Lys Phe Tyr Glu Leu Tyr Glu Leu Asp Ala Glu Leu Gly 295 His Lys Glu Leu Asp Trp Lys Met Thr Met Ser Gly Val Gly Lys Cys 305 310 Arg Gln Val Gly Ile Ser Glu Ser Gly Ile Asp Glu Ala Val Gln Lys 325 Leu Leu Ala Arg Gly Tyr Lys Val Gly Arg Ile Glu Gln Leu Glu Thr 345 Ser Asp Gln Ala Lys Ala Arg Gly Ala Asn Thr Ile Ile Pro Arg Lys 360 Leu Val Gln Val Leu Thr Pro Ser Thr Ala Ser Glu Gly Asn Ile Gly 375 380 370 Pro Asp Ala Val His Leu Leu Ala Ile Lys Glu Ile Lys Met Glu Leu 390 385 Gln Lys Cys Ser Thr Val Tyr Gly Phe Ala Phe Val Asp Cys Ala Ala

410

to the first

Leu	Arg	Phe	Trp 420	Val	Gly	Ser	Ile	Ser 425	Asp	Asp	Ala	Ser	Cys 430	Ala	Ala
Leu	Gly	Ala 435	Leu	Leu	Met	Gln	Val 440	Ser	Pro	Lys	Glu	Val 445	Leu	Tyr	Asp
Ser	Lys 450	Gly	Leu	Ser	Arg	Glu 455	Ala	Gln	Lys	Ala	Leu 460	Arg	Lys	Tyr	Thr
Leu 465	Thr	Gly	Ser	Thr	Ala 470	Val	Gln	Leu	Ala	Pro 475	Val	Pro	Gln	Val	Met 480
Gly	Asp	Thr	Asp	Ala 485	Ala	Gly	Val	Arg	Asn 490	Ile	Ile	Glu	Ser	Asn 495	Gly
Tyr	Phe	Lys	Gly 500	Ser	Ser	Glu	Ser	Trp 505	Asn	Cys	Ala	Val	Asp 510	Gly	Leu
Asn	Glu	Cys 515	Asp	Val	Ala	Leu	Ser 520	Ala	Leu	Gly	Glu	Leu 525	Ile	Asn	His
Leu	Ser 530	Arg	Leu	Lys	Leu	Glu 535	Asp	Val	Leu	Lys	His 540	Gly	Asp	Ile	Phe
Pro 545	Tyr	Gln	Val	Tyr	Arg 550	Gly	Cys	Leu	Arg	Ile 555	Asp	Gly	Gln	Thr	Met 560
Val	Asn	Leu	Glu	Ile 565	Phe	Asn	Asn	Ser	Cys 570	Asp	Gly	Gly	Pro	Ser 575	Gly
Thr	Leu	Tyr	Lys 580	Tyr	Leu	Asp	Asn	Cys 585	Val	Ser	Pro	Thr	Gly 590	Lys	Arg
Leu	Leu	Arg 595	Asn	Trp	Ile	Суѕ	His 600	Pro	Leu	Lys	Asp	Val 605	Glu	Ser	Ile
Asn	Lys 610	Arg	Leu	Asp	Val	Val 615	Glu	Glu	Phe	Thr	Ala 620	Asn	Ser	Glu	Ser
Met 625	Gln	Ile	Thr	Gly	Gln 630	Tyr	Leu	His	Lys	Leu 635	Pro	Asp	Leu	Glu	Arg 640
Leu	Leu	Gly	Arg	Ile 645	Lys	Ser	Ser	Val	Arg 650	Ser	Ser	Ala	Ser	Val 655	Leu
Pro	Ala	Leu	Leu 660	Gly	Lys	Lys	Val	Leu 665	Lys	Gln	Arg	Val	Lys 670	Ala	Phe
Gly	Gln	Ile 675	Val	Lys	Gly	Phe	Arg 680	Ser	Gly	Ile	Asp	Leu 685	Leu	Leu	Ala
Leu	Gln 690	Lys	Glu	Ser	Asn	Met 695	Met	Ser	Leu	Leu	Tyr 700	Lys	Leu	Cys	Lys

Leu 705	Pro	Ile	Leu	Val	Gly 710	Lys	Ser	Gly	Leu	Glu 715	Leu	Phe	Leu	Ser	Gln 720
Phe	Glu	Ala	Ala	Ile 725	Asp	Ser	Asp	Phe	Pro 730	Asn	Tyr	Gln	Asn	Gln 735	Asp
Val.	Thr	Asp	Glu 740	Asn	Ala	Glu	Thr	Leu 745	Thr	Ile	Leu	Ile	Glu 750	Leu	Phe
Ile	Glu	Arg 755	Ala	Thr	Gln	Trp	Ser 760	Glu	Val	Ile	His	Thr 765	Ile	Ser	Cys
Leu	Asp 770	Val	Leu	Arg	Ser	Phe 775	Ala	Ile	Ala	Ala	Ser 780	Leu	Ser	Ala	Gly
Ser 785	Met	Ala	Arg	Pro	Val 790	Ile	Phe	Pro	Glu	Ser 795	Glu	Ala	Thr	Asp	Gln 800
Asn	Gln	Lys	Thr	Lys 805	Gly	Pro	Ile	Leu	Lys 810	Ile	Gln	Gly	Leu	Trp 815	His
Pro	Phe	Ala	Val 820	Ala	Ala	Asp	Gly	Gln 825	Leu	Pro	Val	Pro	Asn 830	Asp	Ile
Leu	Leu	Gly 835	Glu	Ala	Arg	Arg	Ser 840	Ser	Gly	Ser	Ile	His 845	Pro	Arg	Ser
Leu	Leu 850	Leu	Thr	Gly	Pro	Asn 855	Met	Gly	Gly	Lys	Ser 860	Thr	Leu	Leu	Arg
Ala 865	Thr	Cys	Leu	Ala	Val 870	Ile	Phe	Ala	Gln	Leu 875	Gly	Cys	Tyr	Val	Pro 880
Cys	Glu	Ser	Cys	Glu 885	Ile	Ser	Leu	Val	Asp 890	Thr	Ile	Phe	Thr	Arg 895	Leu
Gly	Ala	Ser	Asp 900	Arg	Ile	Met	Thr	Gly 905	Glu	Ser	Thr	Phe	Leu 910	Val	Glu
Cys	Thr	Glu 915		Ala	Ser	Val	Leu 920	Gln	Asn	Ala	Thr	Gln 925	_	Ser	Leu
Val	Ile 930		Asp	Glu	Leu	Gly 935		Gly	Thr	Ser	Thr 940		Asp	Gly	Tyr
Ala 945		Ala	Tyr	Ser	Val 950	Phe	Arg	His	Leu	Val 955		Lys	Val	Gln	Cys 960
Arg	Met	Leu	Phe	Ala 965		His	Tyr	His	Pro		Thr	Lys	Glu	Phe 975	Ala
Ser	His	Pro	Arg 980		Thr	Ser	Lys	His 985		Ala	. Cys	Ala	. Phe 990		Ser

Arg Ser Asp Tyr Gln Pro Arg Gly Cys Asp Gln Asp Leu Val Phe Leu

	995	1000	1005		
Tyr Arg 1010	Leu Thr Glu	Gly Ala Cys Pro	o Glu Ser Tyr Gly 1020	Leu Gln Val	
Ala Leu 1025	-	Ile Pro Asn Gl	n Val Val Glu Thr 1035	Ala Ser Gly 1040	
Ala Ala	Gln Ala Met 1045	Lys Arg Ser Il	e Gly Glu Asn Phe 1050	Lys Ser Ser 1055	
Glu Leu	Arg Ser Glu 1060	Phe Ser Ser Le	ı His Glu Asp Trp 5	Leu Lys Ser 1070	
	Gly Ile Ser 1075	Arg Val Ala Hi 1080	s Asn Asn Ala Pro 1085	Ile Gly Glu	
Asp Asp 1090	Tyr Asp Thr	Leu Phe Cys Le	u Trp His Glu Ile 1100	Lys Ser Ser	
Tyr Cys 1105	Val Pro Lys				
<210>	32				
<211>	24				
<212>	DNA				
<213>	Artifi	cial sequence			
<220>					
<223>		d primer for PC atellite	R amplification o	f ATHGENEA	
<400>	32				
accatgo	ata gcttaaac	tt cttg		24	r
<210>	33				
<211>	22				
<212>	DNA				
<213>	Artifi	cial sequence			
<220>					
<223>		e primer for PC atellite	R amplification o	f ATHGENEA	
<400>	33				
acataac	cac aaataggg	igt gc		22	2

```
<210>
                  34
     <211>
                  18
     <212>
                  DNA
                  Artificial sequence
     <213>
     <220>
                  Forward primer DMCIN-A for PCR on genomic DNA of Arabidopsis
     <223>
                   thaliana ssp. Landsberg erecta "Ler"
     <400>
                   34
                                                                                18
     gaagcgatat tgttcgtg
     <210>
                   35
     <211>
                   18
                   DNA
     <212>
ű
                  Artificial sequence
      <213>
<220>
ű
                   Reverse primer DMCIN-B for PCR on genomic DNA of Arabidopsis
      <223>
thaliana ssp. Landsberg erecta "Ler"
H
      <400>
                   35
The second second second second
                                                                                 18
      agattgcgag aacattcc
                   36
      <210>
                   31
      <211>
      <212>
                   DNA
                   Artificial sequence
      <213>
      <220>
                   Forward primer DMCIN-1 for PCR on genomic DNA of Arabidopsis
      <223>
                   thaliana ssp. Landsberg erecta "Ler"
      <400>
                   36
                                                                                 31
      acgcgtcgac tcagctatga gattactcgt g
      <210>
                    37
                    29
      <211>
                   DNA
      <212>
      <213>
                   Artificial sequence
      <220>
                    Reverse primer DMCIN-2 for PCR on genomic DNA of Arabidopsis
       <223>
                    thaliana ssp. Landsberg erecta "Ler"
       <400>
                    37
```

and the second of the second o

qctctagatt tctcgctcta agactctct

	<210>	38	
	<211>	32	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer DMCIN-3 for PCR on genomic DNA of Arabidopsis	
•	-	thaliana ssp. Landsberg erecta "Ler"	
	<400>	38	
	gctctagagc t	stotottaag taagtgattg at 32	
dograph.	<210>	39	
m) M	<211>	48	
Skuf EN	<212>	DNA	
The state of the s	<213>	Artificial sequence	
i.i	<220>		
1.0 1.0	<223>	Reverse primer DMCIN-4 for PCR on genomic DNA of Arabidopsis	
		thaliana ssp. Landsberg erecta "Ler"	
11	<400>	39	
	teccceggge	tegagagate tecatggttt etteagetet atgaatee 48	;
	<210>	40	
neous Leons	<211>	26	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer DMCla for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"	
	<400>	40	
	acgcgtcgac	gaattcgcaa gtgggg 2	6
*			
	<210>	41	
	<211>	38	
у, .	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Reverse primer DMC1b for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"	
	<400>	41	

and the second of the second o

tccatggaga	a tetecegggt acegatttge ttegaggg	3
<210>	42	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of ATEAT1 SSLP marke Arabidopsis thaliana subspecies	L
<400>	42	
gccactgcg	t gaatgatatg	2
210	42	
<210>	43	
<211>	22 DNA	
<21.2>	DNA Artificial sequence	
<213>	Wifilicial peddence	
<220>		
<223>	Reverse primer for PCR amplification of ATEAT1 SSLP marke Arabidopsis thaliana subspecies	r
<400>	43	
cgaacagco	a acattaattc cc	
<210>	44	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA63 SSLP marker Arabidopsis thaliana subspecies	-
<400>	44	
aaccaaggo	ca cagaagcg	
<210>	45	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA63 SSLP market Arabidopsis thaliana subspecies	r

And the second of the second

	<400>	45	
	acccaagtga t	cgccacc	18
	<210>	46	
	<211>	21	
	<212>	DNA	
!	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA248 SSLP marker Arabidopsis thaliana subspecies	rin
	<400>	46	
Hart Sin and This and Long state Land	taccgaacca a	aaacacaaag g	21
4.j 	<210>	47	
A.	<211>	22	
l.i	<212>	DNA	
7.5 3%	<213>	Artificial sequence	
i.	(213)	MC2220202 Coquation	
=¥.	<220>		
	<223>	Reverse primer for PCR amplification of NGA248 SSLP marke: Arabidopsis thaliana subspecies	r in
	<400>	47	
in the second se	tctgtatctc (ggtgaattet ee	22
	<210>	48	
	<211>	22	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA128 SSLP marke	r in
	(2237	Arabidopsis thaliana subspecies	
*	<400>	48	
*	ggtctgttga	tgtcgtaagt cg	22
	<210>	49	
	<211>	22	
	<211>	DNA	
	<212> <213>	Artificial sequence	
		ALCALLOLUS DOGUCIOC	
	<220>		

<223>	Reverse primer for PCR amplification of NGA128 SSLP marker i Arabidopsis thaliana subspecies	.n
<400>	49	
atcttgaaac	ctttagggag gg 22	2
<210>	50	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA280 SSLP marker in Arabidopsis thaliana subspecies	Ln
<400>	50	
ctgatctcac	ggacaatagt gc 2:	2
<210>	51	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA280 SSLP marker : Arabidopsis thaliana subspecies	in
<400>	51	
ggctccataa	aaagtgcacc 2	0
<210>	52	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA111 SSLP marker Arabidopsis thaliana subspecies	in
<400>	52	
ctccagttgg	aagctaaagg g 2	1
<210>	53	
<211>	21	
<211>	DNA	
<212>	Artificial sequence	
ヘムエンノ	FIL CLEACHER DOG WOLLO	

PCT/EP98/06977 WO 99/19492

38

	<220>		
	<223>	Reverse primer for PCR amplification of NGA111 SSLP marker	in
		Arabidopsis thaliana subspecies	
	< 400 >	53	
3	tgttttttag g	gacaaatggc g	21
f	<210>	54	
	<211>	20	
	<211>	DNA	
		Artificial sequence	
	<213>	Artificial sequence	
	<220>		
berge spik	<223>	Forward primer for PCR amplification of NGA168 SSLP marker	in
w.		Arabidopsis thaliana subspecies	
	<400>	54	
ill'	ccttcacatc	caaaacccac	20
The state of the s			
11 E	<210>	55	
E	<211>	20	
i-i-	<212>	DNA	
	<213>	Artificial sequence	
		•	
4.1	<220>		
the sales by the best of the b	<223>	Reverse primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	55	
	gcacataccc	acaaccagaa	20
	<210>	56	
	<210>	20	
		DNA	
	<212>		
	<213>	Artificial sequence	
*	<220>		
	<223>	Forward primer for PCR amplification of NGA1126 SSLP marke	r
		in Arabidopsis thaliana subspecies	
•			
	<400>	56	
	cgctacgctt	ttcggtaaag	20

the control of the co

	<210>	57
	<211>	20
	<212>	DNA
	<213>	Artificial sequence
	<220>	
	<223>	Reverse primer for PCR amplification of NGA1126 SSLP marker
		in Arabidopsis thaliana subspecies
	<400>	57
	gcacagtcca a	gtcacaacc 20
	<210>	58
	<211>	20
27.00 27.00	<212>	DNA
	<213>	Artificial sequence
71		
	<220>	
	<223>	Forward primer for PCR amplification of NGA361 SSLP marker in
		Arabidopsis thaliana subspecies
and the second s	<400>	58
1		raatttggag 20
	aaagagatga g	raatttggac 20
rt.	<210>	59
2 2 5	<211>	23
ona one	<212>	DNA
	<213>	Artificial sequence
	1110	
	<220>	
	<223>	Reverse primer for PCR amplification of NGA361 SSLP marker in Arabidopsis thaliana subspecies
	<400>	59
	acatatcaat a	atattaaagt agc 23
£	<210>	60
	<211>	18
	<212>	DNA
•	<213>	Artificial sequence
	<220>	
	<223>	Forward primer for PCR amplification of NGA168 SSLP marker in
	-	Arabidopsis thaliana subspecies
	<400>	60

33.4.63		
į,	i	
	Sharp, Heath	9
24,,,,,	Ĭ,	Hilling
	Ē,	
il in.	The part	1111111
	3	
	::	ŧ.
0.	f.	-
the side		
the said of the said	***	100
the said of the said	***	100
the makes thank theme aft'	2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Colonia State March 1811
the makes thank theme aft'	2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Colonia State March 1811
the makes thank theme aft'	***	Colonia State March 1811

	tegtetaetg e	cactgccg	18
	<210>	61	
	<211>	22	
	<212>	DNA	
•	<213>	Artificial sequence	
	000		
,	<220>	Reverse primer for PCR amplification of NGA168 SSLP marke	r in
•	<223>	Arabidopsis thaliana subspecies	
	<400>	61	
	gaggacatgt a	ataggageet eg	22
Secretarion Secre			
. I	<210>	62	
	<211>	20	
î ji	<212>	DNA	
	<213>	Artificial sequence	
111	<220>		
2.	<223>	Forward primer for PCR amplification of AthBIO2 SSLP mark in Arabidopsis thaliana subspecies	cer
Emarkin Emarkin			
	<400>	62	
	tgacctcctc	ttccatggag	20
Topico II	<210>	63	
	<211>	22	
	<211>	DNA	
	<213>	Artificial sequence	
	<220>	n a simon for pan amplification of Atheros SSIP mar	ker
	<223>	Reverse primer for PCR amplification of AthBIO2 SSLP mar in Arabidopsis thaliana subspecies	
	<400>	63	
\$	ttaacagaaa	cccaaagctt tc	22
	<210>	64	
	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		_
	<223>	Forward primer for PCR amplification of AthUBIQUE SSLP m	narker
		in Arabidopsis thaliana subspecies	

```
<400>
             64
                                                                         21
aggcaaatgt ccatttcatt g
             65
<210>
<211>
             20
             DNA
<212>
             Artificial sequence
<213>
<220>
             Reverse primer for PCR amplification of AthUBIQUE SSLP marker
<223>
             in Arabidopsis thaliana subspecies
             65
<400>
acgacatggc agatttctcc
                                                                         20
             66
<210>
<211>
             21
<212>
             DNA
<213>
             Artificial sequence
<220>
             Forward primer for PCR amplification of NGA172 SSLP marker in
<223>
             Arabidopsis thaliana subspecies
<400>
             66
                                                                         21
agetgettee ttatagegte e
<210>
             67
<211>
             19
<212>
             DNA
<213>
             Artificial sequence
<220>
             Reverse primer for PCR amplification of NGA172 SSLP marker in
<223>
             Arabidopsis thaliana subspecies
<400>
                                                                          19
catccgaatg ccattgttc
<210>
             68
<211>
             21
<212>
             DNA
<213>
             Artificial sequence
<220>
```

The second secon

22

20

22

<210> 72 <211> 21 <212> DNA

<213> Artificial sequence

	<220>		
	<223>	Forward primer for PCR amplification of NGA6 SSLP marker in Arabidopsis thaliana subspecies	a
	<400>	72	
	tggatttctt	cctctcttca c	21
r			
	<210>	73	
	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Reverse primer for PCR amplification of NGA6 SSLP marker i	n
The same of the sa	(22)	Arabidopsis thaliana subspecies	
Strong Strong Strong serial Strong St	<400>	73	
4. 14			21
Conjust	atggagaagc	ttacactgat c	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			
	<210>	74	
	<211>	20	
etile Man	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	74	
	aatgttgtcc	tacatacta	20
	0.7.0	75	
	<210> <211>	75 22	
	<211> <212>	DNA	
	<213>	Artificial sequence	
<u>\$</u>	<220>		
	<223>	Reverse primer for PCR amplification of NGA12 SSLP marker	in
		Arabidopsis thaliana subspecies	
ř.			
	<400>	75	
	tgatgctctc	tgaaacaaga gc	22

and the second of the second o

PCT/EP98/06977

44

WO 99/19492

	<210>	76	
	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA8 SSLP marker in	L
		Arabidopsis thaliana subspecies	
	<400>	76	
			٠,
	gagggcaaat	ctttatttcg g	21
	<210>	77	
55 KE	<211>	22	
#555F	<212>	DNA	
	<213>	Artificial sequence	
		*	
	<220>		
T.	<223>	Reverse primer for PCR amplification of NGA8 SSLP marker in	1
IJ		Arabidopsis thaliana subspecies	
Æ	<400>	77	
Will the state of			
	tggctttcgt	ttataaacat cc	22
-			
i i trust	<210>	78	
1 1	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA1107 SSLP market	_
	12007	in Arabidopsis thaliana subspecies	•
		in managers character samplestes	
	<400>	78	
	gcgaaaaaac	aaaaaaatcc a	21
	T.		
y	<210>	79	
	<211>	21	
,	<212>	DNA	
<u> </u>	<213>	Artificial sequence	
	<220>		
	<223>	Reverse primer for PCR amplification of NGA1107 SSLP market	
		in Arabidopsis thaliana subspecies	
	<400>	79	

	cgacgaatcg	acagaattag g	21
	<210>	80	
	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
	(213)	Altificial Sequence	
	<220>		
Í	<223>	Forward primer for PCR amplification of NGA225 SSLP marke: Arabidopsis thaliana subspecies	r in
	<400>	80	
	gaaatccaaa	tcccagagag g	21
engers or significant signifi	-210-	0.1	
	<210>	81	
	<211>	22	
Ti.	<212>	DNA	
the state of the s	<213>	Artificial sequence	
1 145 1 2 3 1 2 3	<220>		
Topic and the second se	<223>	Reverse primer for PCR amplification of NGA225 SSLP marke: Arabidopsis thaliana subspecies	r in
	<400>	81	
Hard Gran Sawa Sawa Sara	tctccccact	agttttgtgt cc	22
o horage	<210>	82	
	<211>	19	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA249 SSLP market Arabidopsis thaliana subspecies	r in
	<400>	82	
	taccgtcaat	ttcatcgcc	19
₹. - 2 ⁸			
	<210>	83	
	<211>	22	
-	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Reverse primer for PCR amplification of NGA249 SSLP marke	r in
	-220	Arabidopsis thaliana subspecies	T T#1

the second second second second

```
<400>
                   83
      ggatccctaa ctgtaaaatc cc
                                                                              22
      <210>
                   84
      <211>
                  22
      <212>
                  DNA
      <213>
                  Artificial sequence
      <220>
                  Forward primer for PCR amplification of CA72 SSLP marker in
      <223>
                  Arabidopsis thaliana subspecies
      <400>
                  84
     aatcccagta accaaacaca ca
                                                                              22
<210>
                  85
ħ
     <211>
                  20
<212>
                  DNA
     <213>
                  Artificial sequence
.S
     <220>
     <223>
                  Reverse primer for PCR amplification of CA72 SSLP marker in
Arabidopsis thaliana subspecies
     <400>
                  85
     cccagtctaa ccacgaccac
                                                                              20
     <210>
                  86
     <211>
                  20
     <212>
                  DNA
     <213>
                  Artificial sequence
     <220>
                  Forward primer for PCR amplification of NGA151 SSLP marker in
     <223>
                  Arabidopsis thaliana subspecies
     <400>
                  86
     gttttgggaa gttttgctgg
                                                                              20
     <210>
                  87
     <211>
                  24
     <212>
                  DNA
     <213>
                  Artificial sequence
```

The state of the s

<220>

PCT/EP98/06977

<213>

Artificial sequence

	<223>	Arabidopsis thaliana subspecies	711
	<400>	87	
	cagtctaaaa	gcgagagtat gatg 2	4
)			
	<210>	88	
i	<211>	22	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA106 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	88	
The first way family from most facility for the family family from the family f	gttatggagt	ttctagggca cg 2	22
	<210>	89	
	<211>	20	
	<212>	DNA	
Marie	<213>	Artificial sequence	
£	<220>		
	<223>	Reverse primer for PCR amplification of NGA106 SSLP marker Arabidopsis thaliana subspecies	in
Testa signi	<400>	89	
	tgccccattt	tgttcttctc	20
	<210>	90	
	<211>	20	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA139 SSLP marker	in
7.7		Arabidopsis thaliana subspecies	
	<400>	90	
•	agagetacea	gatccgatgg	20
	<210>	91	
	<211>	21	
	<212>	DNA	

WO 99/19492 PCT/EP98/06977

48

	<220>		
	<223>	Reverse primer for PCR amplification of NGA139 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	91	
ı	ggtttcgttt	cactatecag g	21
ŧ	0.7.0		
	<210>	92	
	<211>	22	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Forward primer for PCR amplification of NGA76 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	92	
Park from the from th	ggagaaaatg	tcactctcca cc	22
in Tipe M. br>M. M. M. M. M. M. M.	<210>	93	
	<211>	20	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Reverse primer for PCR amplification of NGA76 SSLP marker Arabidopsis thaliana subspecies	in
	<400>	93	
	aggcatggga	gacatttacg	20
	<210>	94	
	<211>	20	
	<212>	DNA	
	<213>	Artificial sequence	
: ,	<220>		
	<223>	Forward primer for PCR amplification of ATHSO191 SSLP mark	
		in Arabidopsis thaliana subspecies	-er
\$		<u>-</u>	
	<400>	94	
	ctccaccaat	catgcaaatg	20

49

	<210>	95	
	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
)	<220>		
	<223>	Reverse primer for PCR amplification of ATHSO191 SSLP mar	ker
*		in Arabidopsis thaliana subspecies	
	<400>	95	
	tgatgttgat	ggagatggtc a	21
Name of the last o	<210>	96	
Marie .	<211>	22	
123 123	<212>	DNA	
e i	<213>	Artificial sequence	
	-220		
	<220> <223>	Daming makes for paper 2151 at a special and	
Manual Control of the	<223>	Forward primer for PCR amplification of NGA129 SSLP marke Arabidopsis thaliana subspecies	rin
1.11	<400>	96	
	tcaggaggaa	ctaaagtgag gg	22
Logi Jerz	<210>	97	
£	<211>	22	
	<212>	DNA	
	<213>	Artificial sequence	
	<220>		
	<223>	Payarsa primar for DCD amplification of NGA120 GGLD ampli-	
	12237	Reverse primer for PCR amplification of NGA129 SSLP marke Arabidopsis thaliana subspecies	r in
	<400>	97	
	cacactgaag	atggtcttga gg	22
t, e	<210>	98	
	<210>		
	<211>	8062 DNA	
<u>k</u>	<213>		
	~2137	Arabidopsis thaliana ecotype Columbia	
	<220>		
	<223>	Genomic DNA sequence of AtMSH6	
	<400>	97	
	ttttttggtt	gctaacaata aaggtatacg gttttatgtc atcaatataa ctatatataa	60

aagaaatgaa agatatatat tgttttttca tttatcaaac aaaacaacaa gactttttt 120 ttacttttta cattggtcaa caaaatacaa gataaacgac atcgtttaat catttcccaa 180 ttttacccct aagtttaaca cctagaacct tctccatctt cgcaagcaca gcctgattag 240 gaacagettt accattetea tatteetgaa etacetgagt eeteteattg atetgttteg 300 ccaaatccgc ttgtgacatc ttcttctcca atctcgcttt ctgtatcatc aacctcacct 360 ctgctttcac acgatccatc gccgcaggct ctgtttcttc ttccagcttc ttcgtgttaa 420 tcaccggaac cgccgtagat ttcccctttt tgttcgaacc ggcatcgaat ttcttaaccg 480 tttgaaccgc gacaccgttt ctcagagctg cgttaaccgc tttcggatcg cgtaggtctt 540 ggctcttttg ttttgatttg tggagaacta ctggttccca gtcttgtgtt actgctcctg 600 ggtatctgct cggcatcgtc gatgaattga gagaaaggaa caacgcgaaa attttattaa 660 totgagtttt gaaattgaga aacgatgaag atgaagaatg tigttgagag gattgtgata 720 tttatatata cgaagattgg tttctggaga attcgatcat ctttttctcc attttcgtct 780 ctggaacgtt cttagagatg attgacgacg tgtcattatc tgatttgcag ttaaccaatg 840 ctttttgggt tggattcgtg gtacaccata ttatccgatt tggctcaatg gttttatata 900 aatttggttt teggtteggt tatgagttat eattaaaatt aagetaacea aaaatttteg 960 taaaatttat tteggtttea atteggatee ettaetteea gaacegaatt attegaaace 1020 ggggttagcc gaaccgaata ccaatgcctg attgactcgt tggctagaaa gatccaacgg 1080 tatacaataa tagaacataa atcggacggt catcaaagcc tcaaagagtg aacagtcaac 1140 aaaaaaagtt gagccctgag gagtatcgtt tccgccattt ctacgacgca aggcgaaaat 1200 ttttggcgcc aatctttccc ccctttcgaa ttctctcagc tcaaaacatc gtttctctct 1260 cactetetet cacaatteca aaaaatgeag egecagagat egattttgte tttettecaa 1320 aaacccacgg cggcgactac gaagggtttg gtttccggcg atgctgctag cggcgggggc 1380 ggcagcggag accacgattt aatgtgaagg aaggggatgc taaaggcgac gcttctgtac 1440 gttttgctgt ttcgaaatct gtcgatgagg ttagaggaac ggatactcca ccggagaagg 1500 ttccgcgtcg tgtcctgccg tctggattta agccggctga atccgccggt gatgcttcgt 1560 ccctgttctc caatattatg cataagtttg taaaagtcga tgatcgagat tgttctggag 1620 agaggtacta atettegatt etettaattt tgttatettt agetggaaga agaagatteg 1680

and a substance of the

tgtaatttgt tgtattcgtt ggagagattc tgattactgc attggatcgt tgtttacaaa 1740 ttttcaggag ccgagaagat gttgttccgc tgaatgattc atctctatgt atgaaggcta 1800 atgatgttat teeteaattt egtteeaata atggtaaaae teaagaaaga aaccatgett 1860 ttagtttcag tgggagagct gaacttagat cagtagaaga tataggagta gatggcgatg 1920 ttcctggtcc agaaacacca gggatgcgtc cacgtgcttc tcgcttgaag cgagttctgg 1980 aggatgaaat gacttttaag gaggataagg ttcctgtatt ggactctaac aaaaggctga 2040 aaatgctcca ggatccggtt tgtggagaga agaaagaagt aaacgaagga accaaatttg 2100 aatggettga gtettetega atcagggatg ceaatagaag aegteetgat gateceettt 2160 acgatagaaa gaccttacac ataccacctg atgttttcaa gaaaatgtct gcatcacaaa 2220 agcaatattg gagtgttaag agtgaatata tggacattgt gcttttcttt aaagtggtta 2280 gtaactatta atctagtgtt caatccattt cctcaatgtg atttgttcac ttacatctgt 2340 ttacgttatg ctcttctcag gggaaatttt atgagctgta tgagctagat gcggaattag 2400 gtcacaagga gcttgactgg aagatgacca tgagtggtgt gggaaaatgc agacaggtaa 2460 attagttgaa acaactggcc tgcttgaatt attgtgtcta taaattttga caccaccttt 2520 tgtttcaggt tggtatctct gaaagtggga tagatgaggc agtgcaaaag ctattagctc 2580 gtgggtaagg gaaccatcat actttatgga attcgtttac tgctacttcg gctaggattt 2640 aagaaatgga aatcacttca agcatcatta gttaggatcc tgagaactca ggatgttttc 2700 ttattcgtta tataataagt cttttcatca aggagtaaca aacaaaactt gcacaatatt 2760 tgtgtgctca ctggcaaggc atatataccc agctaacctt tgctagttca ctgtagtaac 2820 agttacggat aatatatgtt tacttgtatg tggtaccctc attttgtctc tcatggaggc 2880 tttcaagcet tgtgttgaaa etggatagtt acatatgett ecaacagaaa etagcatgea 2940 gattcatatg ctttcctatt ctactaatta tgtattgaca cactcgttgt ttcttttgaa 3000 agatataaag ttggacgaat cgagcagcta gaaacatctg accaagcaaa agccagaggt 3060 gctaatactg taagttttct tggataggtc aaggagagtg ttgcagactg tttttgatca 3120 tttctttttc tgtacattac tttcatgctg taattaactc aatggctatt ctggtctgat 3180 tatcagataa ttccaaggaa gctagttcag gtattaactc catcaacagc aagcgaggga 3240 aacatcgggc ctgatgccgt ccatcttctt gctataaaag aggtttgtta tttacttatt 3300

باللانتين ببالبني بي

3360 tatettatea tgtteagtte atecaagtee tgaaaaatta caetettett taccaatett 3420 ccatcaagct gtgtaaagga tttggaatta gaaaatcatt atttgatgct ttgttttata tgcaagaggt tcccttgaaa agatctgitt aagattcttt gcacttgaaa aattcaatct 3480 3540 ttttaagtga atcccctact ttcttacaat gatcatagtc tgcaattgca tgtcaagtaa 3600 tateatteet tgttaetgea tecceetett tettaatgae cattgtetat gttgtgtttg tctcgtgtgc tggagaaaat gatagctgat ccaagctgta cattatcatg attaagtagc 3660 3720 tgctcaggaa ttgcctttgg ttacattgcc taatggtttg atgtcaattt ttcttctgaa tetttatttt agateaaaat ggagetaeaa aagtgtteaa etgtgtatgg atttgetttt 3780 3840 gttgactgtg ctgccttgag gttttgggtt gggtccatca gcgatgatgc atcatgtgct getettggag egitattgat geaggiaage aagigtatte igiatettat gigtaecaig 3900 tgacttcctg tgcatatatt tgggttgcag gaactaattc tgaatcacca tttggtatgt 3960 4020 tttttccagg tttctccaaa ggaagtgtta tatgacagta aaggtaaact gcttgtatcg 4080 ccagttgttt tgttaaacag aatttaaggt aaatgacact ggttaattta aagtgcatac 4140 atgttgaaat attgcagggc tatcaagaga agcacaaaag gctctaagga aatatacgtt 4200 gacaggtace attteagtag geaagetaae tgacaattta acegeteace gaatgatagg 4260 tetettaaae attgetaatg tagatgatgt ttatgtttea atetaatagg gtetaeggeg 4320 gtacagttgg ctccagtacc acaagtaatg ggggatacag atgctgctgg agttagaaat ataatagaat ctaacggata ctttaaaggt tcttctgaat catggaactg tgctgttgat 4380 4440 ggtctaaatg aatgtgatgt tgcccttagt gctcttggag agctaattaa tcatctgtct 4500 aggctaaagg tgtgttggct tgtttagttt ttgcttttca caaattaagc aaaggaactt 4560 ttcataactt acagtttcta tctacttgca gctagaagat gtacttaagc atggggatat ttttccatac caagtttaca ggggttgtct cagaattgat ggccagacga tggtaaatct 4620 4680 tgagatattt aacaatagct gtgatggtgg tccttcaggc aagtgcatat ttcttttttg ataacttcaa ctagagggca gacatagaag gaaaaattct aatacttcgt acggatctcc 4740 agtaagtaat agccgatttt tgtttaccta tgtagggacc ttgtacaaat atcttgataa 4800 ctgtgttagt ccaactggta agcgactctt aaggaattgg atctgccatc cactcaaaga 4860 tgtagaaagc atcaataaac ggcttgatgt agttgaagaa ttcacggcaa actcagaaag 4920

.*

4980 tatgcaaatc actggccagt atctccacaa acttccagac ttagaaagac tgctcggacg 5040 catcaagtet agegttegat catcageete tgtgttgeet getettetgg ggaaaaaagt gctgaaacaa cgagtaagta tcaatcacaa gttttctgag taatgccttc catgagtagt 5100 5160 ataggactaa aacattacgg gtctagctaa agactgttct ccttcttttg caatgtctgg 5220 ttattcatta catttctctt aacttattgc attgcaggtt aaagcatttg ggcaaattgt 5280 gaaagggttc agaagtggaa ttgatctgtt gttggctcta cagaaggaat caaatatgat gagtttgctt tataaactct gtaaacttcc tatattagta ggaaaaagcg ggctagagtt 5340 5400 atttetttet caattegaag cagecataga tagegaettt ceaaattate aggtgeecat ctatctttca tactttacaa caaaatgtct gtcactactc aaagcaatgc atatggctta 5460 5520 gatotoaact cacaccocga ggatoctaaa gggatttget tittaittoot aatgittitig 5580 gatggtttga tttatttcta acttgaactt attaatcttg taccagaacc aagatgtgac agatgaaaac gctgaaactc tcacaatact tatcgaactt tttatcgaaa gagcaactca 5640 5700 atggtctgag gtcattcaca ccataagctg cctagatgtc ctgagatctt ttgcaatcgc 5760 agcaagtete tetgetggaa geatggeeag geetgttatt ttteeegaat cagaagetae 5820 agatcagaat cagaaaacaa aagggccaat acttaaaatc caaggactat ggcatccatt 5880 tgcagttgca gccgatggtc aattgcctgt tccgaatgat atactccttg gcgaggctag 5940 aagaagcagt ggcagcattc atcctcggtc attgttactg acgggaccaa acatgggcgg 6000 aaaatcaact cttcttcgtg caacatgtct ggccgttatc tttgcccaag tttgtatact 6060 cgttagataa ttactctatt ctttgcaatc agttcttcaa catgaataat aaattctgtt 6120 ttetgtetge agettggetg etaegtgeeg tgtgagtett gegaaatete eetegtggat 6180 actatettea caaggettgg egeatetgat agaateatga caggagagag taagttttgt 6240 teteaaaata eeaatteete gaactattta eteagatttt gtetgattgg acaaggtggt 6300 tttgcttttt tttaggtacc tttttggtag aatgcactga gacagcgtca gttcttcaga atgcaactca ggattcacta gtaatccttg acgaactggg cagaggaact agtactttcg 6360 6420 atggatacgc cattgcatac teggtaacct getettetee tteaacttat acttgttgat 6480 caacaaaaac atgcaattca ttttgctgaa acttattgat ttatatcagg tttttcgtca 6540 cctggtagag aaagttcaat gtcggatgct ctttgcaaca cattaccacc ctctcaccaa

ggaattcgcg	tctcacccac	gtgtcacctc	gaaacacatg	gcttgcgcat	tcaaatcaag	6600
atctgattat	caaccacgtg	gttgtgatca	agacctagtg	ttcttgtacc	gtttaaccga	6660
gggagcttgt	cctgagagct	acggacttca	agtggcactc	atggctggaa	taccaaacca	6720
agtggttgaa	acagcatcag	gtgctgctca	agccatgaag	agatcaattg	gggaaaactt	6780
caagtcaagt	gagctaagat	ctgagttctc	aagtctgcat	gaagactggc	tcaagtcatt	6840
ggtgggtatt	tctcgagtcg	cccacaacaa	tgcccccatt	ggcgaagatg	actacgacac	6900
tttgttttgc	ttatggcatg	agatcaaatc	ctcttactgt	gttcccaaat	aaatggctat	6960
gacataacac	tatctgaagc	tegttaagte	ttttgcttct	ctgatgttta	ttcctcttaa	7020
aaaatgctta	tatatcaaaa	aattgtttcc	tcgattataa	caagattata	tatgtatctg	7080
tcggtttagc	tatggtatat	aatatatgta	tgttcatgag	attggtcaag	agaaatactc	7140
acaaacagta	tattaagaag	gaaatatgtt	tatgcattaa	tttaagtttc	aagataaact	7200
gcaaataacc	tcgactaaag	ttgcaaagac	caaacacaaa	ttacaaaact	tataagactt	7260
aagttctgaa	ttccctaaaa	ccaaaaaaaa	aaacagaaca	tattttgttg	catctacaaa	7320
caacacaaac	ctacatagtt	tataacttac	tcatcactga	gattaacatc	agaatcattc	7380
tccatttctt	catcttcact	ctcatcatca	tcaccaccac	catgatgatt	ctcctcct	7440
tcacgtaacc	tagcaatctc	actctgagct	ctatcaacaa	tetgettett	ctgcaactcc	7500
aaatctctct	gaaaatcagc	tctcatcttc	tccaactcct	tcatttgctc	tttcttactc	7560
ttctccatct	tctcataaac	cttcccaaac	ctctcaacag	aatccgccaa	catcttatac	7620
gaagcagcgt	cattaacctt	cttcctctcg	tactcaacct	catcatcctc	atcctcctcc	7680
tcttcagaat	caccaggact	atccatcatc	tcatcaaacc	cattagactt	atctaaataa	7740
accttagtgt	tcataaacac	aaactcacct	gaatcaacac	cacaagctaa	acctaaatcc	7800
gacttgggcg	aaacacaaag	caacatatcc	aacttattga	aaaacgacca	tttacttgaa	7860
cctaaacctg	atttctcaac	cttaatcttc	tcttttctat	acttcctctt	caagtcatca	7920
atcattctcc	tacattgcgt	ctcagatttc	tccatcctta	gctcctcact	cactttctca	7980
gctacttcat	tccaatcctc	gttcctcaaa	ctccttctac	ccaattgcaa	aaacctatct	8040
ccccaaactt	caagcaacac	aa				8062

BAKER BOTTS L.L.P. FILE NO.: A33153-PCT-USA-072667.0128

COMBINED DECLARATION AND POWER OF ATTORNEY

(Original, Design, National Stage of PCT, Divisional, Continuation or C-I-P Application)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHODS FOR OBTAINING PLANT VARIETIES

This declaration is of the following type:
[] original
[] design
[X] national stage of PCT.
Fil distributed
[] continuation
[] continuation [[] continuation-in-part (C-I-P)
the specification of which: (complete (a), (b), or (c))
(a) [] is attached hereto.
(b) [X] was filed on April 10, 2000 as Application Serial No. 09/529,239 and was amended on (if applicable)
(c) [X] was described and claimed in PCT International Application No. PCT/EP98/06977 filed on Octobe
10, 1997 and was amended on (if applicable).
Acknowledgement of Review of Papers and Duty of Candor I hereby state that I have reviewed and understand the contents of the above identified specification
Acknowledgement of Review of Papers and Duty of Candor
I hereby state that I have reviewed and understand the contents of the above identified specification
including the claims, as amended by any amendment referred to above.
I acknowledge the duty to disclose information which is material to the patentability of the subject matter
claimed in this application in accordance with Title 37, Code of Federal Regulations § 1.56.
[] In compliance with this duty there is attached an information disclosure statement. 37 CFR 1.98.
Priority Claim
I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) of any foreign
application(s) for patent or inventor's certificate or of any PCT International Application(s) designating at least on

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) of any foreign application(s) for patent or inventor's certificate or of any PCT International Application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT International Application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application on which priority is claimed

(complete (d) or (e))

- (d) [] no such applications have been filed.
- (e) [X] such applications have been filed as follows:

BAKER BOTTS L.L.P.

FILE NO.: A33153-PCT-USA-072667.0128

COUNTRY	APPLICATION NO.	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
Australia	PO9745	October 10, 1997	-	[X]YES NO[]
				[] YES NO []
				[] YES NO []
L FOREIGN APPL	ICATION[S], IF ANY, FILED MORE TH	AN 12 MONTHS (6 MONTHS FOR DESIGN) PRIC	OR TO SAID APPLICATION	
	-			[]YES NO []
				[]YES NO []
				[]YES NO []

Claim for Benefit of Prior U.S. Provisional Application(s)

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below:

Provisional Application Number	Filing Date	
7		

Claim for Benefit of Earlier U.S./PCT Application(s) under 35 U.S.C. 120

(complete this part only if this is a divisional, continuation or C-I-P application)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior application(s) in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose information as defined in Title 37, Code of Federal Regulations, § 1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned

Power of Attorney

As a named inventor, I hereby appoint Dana M. Raymond, Reg. No. 18,540; Frederick C. Carver, Reg. No. 17,021; Francis J. Hone, Reg. No. 18,662; Joseph D. Garon, Reg. No. 20,420; Arthur S. Tenser, Reg. No. 18,839; Ronald B. Hildreth, Reg. No. 19,498; Thomas R. Nesbitt, Jr., Reg. No. 22,075; Robert Neuner, Reg. No. 24,316; Richard G. Berkley, Reg. No. 25,465; Richard S. Clark, Reg. No. 26,154; Bradley B. Geist, Reg. No. 27,551; James J. Maune, Reg. No. 26,946; John D. Murnane, Reg. No. 29,836; Henry Tang, Reg. No. 29,705; Robert C. Scheinfeld, Reg. No. 31,300; John A. Fogarty, Jr., Reg. No. 22,348; Louis S. Sorell, Reg. No. 32,439; Rochelle K. Seide Reg. No. 32,300; Gary M. Butter, Reg. No. 33,841; Marta E. Delsignore, Reg. No. 32,689; Lisa B. Kole, Reg. No. 35,225 and Janet M. MacLeod, Reg. No. 35,263 of the firm of BAKER BOTTS L.L.P., with offices at 30 Rockefeller Plaza, New York, New York 10112, as attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith

SEND CORRESPONDENCE TO: BAKER BOTTS L.L.P. 30 ROCKEFELLER PLAZA, NEW YORK, N.Y. 10112- CUSTOMER NUMBER: 21003	DIRECT TELEPHONE CALLS TO: BAKER BOTTS L.L.P. (212) 705-5000
---	---

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge

NY02:270141.1 -2-

BAKER BOTTS L.L.P.

FILE NO.: A33153-PCT-USA-072667.0128

that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

	LAST NAME	FIRST NAME	MIDDLE NAME	
FULL NAME OF SOLE OR FIRST INVENTOR			MIDDLE NAME	
	DOUTRIAUX	MARIE-PASCALE		
RESIDENCE & CITIZENSHIP	1	STATE or FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	Saulx les Chartreux	FRANCE FKX	FRANCE	
POST OFFICE	POST OFFICE ADDRESS	CITY	STATE or COUNTRY	ZIP CODE
ADDRESS	64, route de Villebon	Saulx les Chartreux	FRANCE	F-91160
DATE	SIGNATURE OF INVENTOR			
f^{\pm}	1/1 _			
x b ocholer 200	ar all			
FÜLL NAME OF SECOND	EAST NAME	FIRST NAME	MIDDLE NAME	
JOINT INVENTOR, IF ANY	BETZNEK	ANDREAS	STEFAN	
RESIDENCE & CITIZENSHIP	CITY	STATE or FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	PAGE	AUSTRALIA	GERMANY	
POST OFFICE				I
POST OFFICE	POST OFFICE ADDRESS	CITY	STATE or COUNTRY	ZIP CODE
ADDRESS	40 Dallachy Place	PAGE FULL	AUSTRALIA	Act 2614
15 September 200	SIGNATURE OF DIVENTOR			
FULL NAME OF THIRD	L'AST NAME	FIRST NAME	MIDDLE NAME	
IOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP	FREYSSINET	GEORGES		
	The state of the s	STATE or FOREIGN COUNTRY	COUNTRY OF CITIZEN	CLITD
RESIDENCE & CITIZENSHIP		, , , ,		
¥ E	Saint Cyr au Mont d'Or	FRANCE TOL	FRANCE	
POST OFFICE	POST OFFICE ADDRESS	CITY	STATE or COUNTRY	ZIP CODE
ADDRESS	21 rue de Nervieux	Saint Cyr au Mont d'Or	FRANCE	F-69450
DATE	CICNIA TUBE OF INIVENTOR		ı	
39 Sylvembre 2000	366			
FULL NAME OF FOURTH	LAST NAME	FIRST NAME	MIDDLE NAME	
JOINT INVENTOR, IF ANY	PEREZ _	PASCAL		
RESIDENCE & CITIZENSHIP	CITY	STATE or FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
Hydr				
Trans.	WADENNIES	ERANCE #1/	I ER A NICE	
and a second	VARENNES	FRANCE TO	FRANCE	
POST OFFICE	POST OFFICE ADDRESS	CITY	STATE or COUNTRY	ZIP CODE
		110/	1	ZIP CODE F-63450
ADDRESS	POST OFFICE ADDRESS 17, chemin de la Pradelle	CITY	STATE or COUNTRY	1
ADDRESS	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR	CITY	STATE or COUNTRY	1
ADDRESS DATE 28 MA Augusta	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR	Varennes	STATE of COUNTRY FRANCE	1
ADDRESS DATE 28 Physical Parties 12 Physical Parties 12 Physical Parties 12 Physical Physical Parties 12 Physical Physic	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR	CITY	STATE or COUNTRY	1
ADDRESS DATE 28 HA Augus 12 FULL NAME OF FIFTH	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR	Varennes	STATE OF COUNTRY FRANCE MIDDLE NAME	1
ADDRESS DATE 28 Hayel FULL NAME OF FIFTH JOINT INVENTOR, IF ANY	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME	Varennes	STATE OF COUNTRY FRANCE MIDDLE NAME	F-63450
ADDRESS DATE 28 HAPPEN 12 FULL NAME OF FIFTH JOINT INVENTOR, IF ANY	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME	Varennes FIRST NAME	STATE OF COUNTRY FRANCE MIDDLE NAME	F-63450
DATE 28 Augus 20 FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME	Varennes FIRST NAME	STATE OF COUNTRY FRANCE MIDDLE NAME	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN	F-63450
DATE 28 Augus 28 FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE FULL NAME OF SIXTH	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY	F-63450
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE FULL NAME OF SIXTH JOINT INVENTOR, IF ANY	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY FIRST NAME	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY	F-63450 SHIP ZIP CODE
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY POST OFFICE ADDRESS DATE FULL NAME OF SIXTH JOINT INVENTOR, IF ANY	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY	F-63450 SHIP ZIP CODE
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY POST OFFICE ADDRESS DATE FULL NAME OF SIXTH JOINT INVENTOR, IF ANY	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY FIRST NAME	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY	F-63450 SHIP ZIP CODE
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE FULL NAME OF SIXTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY FIRST NAME	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY	F-63450 SHIP ZIP CODE
DATE 28 Augus 28 FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE FULL NAME OF SIXTH	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME CITY	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY MIDDLE NAME COUNTRY OF CITIZEN	F-63450 SHIP ZIP CODE
ADDRESS DATE 28 August FULL NAME OF FIFTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP POST OFFICE ADDRESS DATE FULL NAME OF SIXTH JOINT INVENTOR, IF ANY RESIDENCE & CITIZENSHIP	POST OFFICE ADDRESS 17, chemin de la Pradelle SIGNATURE OF INVENTOR LAST NAME CITY POST OFFICE ADDRESS SIGNATURE OF INVENTOR LAST NAME CITY	CITY Varennes FIRST NAME STATE or FOREIGN COUNTRY CITY FIRST NAME STATE or FOREIGN COUNTRY	STATE OF COUNTRY FRANCE MIDDLE NAME COUNTRY OF CITIZEN STATE OF COUNTRY MIDDLE NAME COUNTRY OF CITIZEN	F-63450 SHIP ZIP CODE